



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number : **0 540 334 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number : 92309924.6

(51) Int. Cl.⁵ : C07D 209/46, A61K 31/395

(22) Date of filing : 29.10.92

(30) Priority : 29.10.91 US 784484
15.01.92 US 821116

(43) Date of publication of application :
05.05.93 Bulletin 93/18

(84) Designated Contracting States :
CH DE FR GB IT LI NL


(71) Applicant : MERCK & CO. INC.
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900 (US)

(72) Inventor : Hartman, George D.
1529 Tennis Circle
Lansdale, PA 19446 (US)
Inventor : Egbertson, Melissa
1232 Lois Road
Ambler, PA 19002 (US)
Inventor : Turchi, Laura M.
214 Croft Ridge Drive
Broomall, PA 19008 (US)
Inventor : Birchenough, Laura A.
2058 Maple Avenue, Apt. E2-6
Hatfield, PA 19440 (US)


(74) Representative : Barrett-Major, Julie Diane et al
Merck & Co., Inc. European Patent
Department Terlings Park Eastwick Road
Harlow Essex CM20 2QR (GB)

(54) Fibrinogen receptor antagonists.

(57) Fibrinogen receptor antagonists of the formula :



are disclosed for use in inhibiting the binding of fibrinogen to blood platelets and for inhibiting the aggregation of blood platelets.
wherein G is :



FIELD OF THE INVENTION

This invention relates to the discovery of fibrinogen receptor antagonists of Formula I for use in inhibiting the binding of fibrinogen to blood platelets and inhibiting the aggregation of blood platelets when administered to mammals, preferably humans.

BACKGROUND OF THE INVENTION

The interaction of platelets with the coagulation and fibrinolytic systems in the maintenance of hemostasis may become pathogenic, requiring prevention and treatment. The fibrinogen receptor antagonists of Formula I are useful in treating various diseases related to platelet aggregation and fibrin formation.

An interest in platelet inhibitors has reemerged as a result of a better understanding of the role of platelets and thrombosis in the pathogenesis of vascular disease, including unstable angina, acute myocardial infarction and stroke.

Platelets are cell-like anucleated fragments, found in the blood of all mammals which participate in blood coagulation. Fibrinogen is a glycoprotein present as a normal component of blood plasma. Fibrinogen participates in platelet aggregation and fibrin formation in the blood clotting mechanism. Platelets are deposited at sites of vascular injury where multiple physiological agonists act to initiate platelet aggregation culminating in the formation of a platelet plug to minimize blood loss. If the platelet plug occurs in the lumen of a blood vessel, normal blood flow is impaired.


Platelet membrane receptors are essential in the process of platelet adhesion and aggregation. Interaction of fibrinogen with a receptor on the platelet membrane complex IIb/IIIa is known to be essential for normal platelet function.

Zimmerman et al., U.S. Patent No. 4,683,291, describes peptides having utility in the study of fibrinogen-platelet, platelet-platelet, and cell-cell interactions. The peptides are described as having utility where it is desirable to retard or prevent formation of a thrombus or clot in the blood. The general formula for the peptides is:




where Ch and Cx are sequences of amino acids.

Pierschbacher et al., U.S. Patent No. 4,589,881, describes the sequence of an 11.5 kDal polypeptide fragment of fibronectin which embodies the cell-attachment-promoting activity of fibronectin. A specifically described fragment is:



Ruoslahti et al., U.S. Patent No. 4,614,517, describes tetrapeptides which alter cell-attachment activity of cells to various substrates. The peptides are stated to "consist essentially of" the following sequence:



wherein X is H or one or more amino acids and Y is OH or one or more amino acids. Figure 1 lists the polypeptides that were synthesized by Ruoslahti et al. in "determining the smallest peptide exhibiting cell attachment activity". Ruoslahti et al., U.S. Patent No. 4,578,079, describes similar tetrapeptides having Ser substituted with Thr or Cys.


Pierschbacher et al., *Proc. Natl. Acad. Sci. USA*, Vol. 81, pp.5985-5988, October, 1984, describe variants of the cell recognition site of fibronectin that retain attachment-promoting activity. Pierschbacher et. al. further assayed the cell attachment-promoting activities of a number of structures closely resembling the Arg-Gly-Asp-Ser peptide, and found "that the arginine, glycine, and aspartate residues cannot be replaced even with closely related amino acids, but that several amino acids can replace serine without loss of activity."

Ruoslahti et al., *Science*, Vol. 238, pp. 491-497, October 23, 1987, discuss cell adhesion proteins. They specifically state that "elucidation of the amino acid sequence of the cell-attachment domain in fibronectin and its duplication with synthetic peptides establish the sequence Arg-Gly-Asp (RGD) as the essential structure

recognized by cells in fibronectin."

Cheresh, Proc. Natl. Acad. Sci. USA, Vol. 84, pp. 6471-6475, September 1987, describes the Arg-Gly-Asp-directed adhesion receptor involved in attachment to fibrinogen and the von Willebrand Factor.

Adams et al., U. S. Patent No. 4,857,508, describes tetrapeptides which inhibit platelet aggregation and the formation of a thrombus. The tetrapeptides have the formula:



wherein X can be



or Ac-Arg, wherein Z = H, NH₂, or NH-Acyl and n=1-4, and wherein Y can be Tyr-NH₂, Phe-NH₂ or a group of a specifically defined formula.

Tjoeng et al., EP 352,249, describe platelet aggregation inhibitors which antagonize interactions between fibrinogen and/or extracellular matrix proteins and the platelet gpIIb/IIIa receptor, including 8-guanido-octanoyl-Asp-2-(4-methoxy-phenyl)ethyl amide.

Alig et al., EP 372,486, describe N-aryl beta-amino acids which inhibit fibrinogen, fibronectin and von Will-ebrand factor to the blood platelet fibrinogen receptor (glyco-protein IIb/IIIa).

Alig et al., EP 381,033, describe di-aryl or heteroaryl substituted alkanolic acid derivatives of a defined formula which inhibit binding of proteins to their specific receptors on cell surfaces, including fibrinogen.

Alig et al., EP 384,362, describe glycine peptides of a specified formula containing an amidine group which inhibit binding of fibrinogen to platelet fibrinogen receptors.


Horwell et al., EP 405,537, describe N-substituted cycloalkyl and polycycloalkyl alpha-substituted Trp-Phe- and phenethylamine derivatives which are useful for treating obesity, hypersecretion of gastric acid in the gut, gastrin-dependent tumors, or as antipsychotics.

It is an object of the present invention to provide fibrinogen receptor antagonists for use in inhibiting the binding of fibrinogen to blood platelets and inhibiting the aggregation of blood platelets. Another aspect of the present invention is to provide novel fibrinogen receptor antagonist compounds. Other objects of the present invention are to provide methods of inhibiting the binding of fibrinogen to blood platelets and inhibiting the aggregation of blood platelets, through the administration of novel fibrinogen receptor antagonist compounds.


The above and other objects are accomplished by the present invention in the manner described below.

SUMMARY OF THE INVENTION

The present invention provides fibrinogen receptor antagonist compounds of the formula:




wherein G is




for use in inhibiting the binding of fibrinogen to blood platelets and for inhibiting the aggregation of blood platelets. The above-mentioned compounds can be used in a method of acting upon a fibrinogen receptor which comprises administering a therapeutically effective but non-toxic amount of such compound to a mammal, preferably a human. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, dispersed therein, an effective but non-toxic amount of such compound is another feature of this invention.

DETAILED DESCRIPTION OF THE INVENTION

Fibrinogen receptor antagonist compounds of Formula I are useful in a method of inhibiting the binding of fibrinogen to blood platelets and for inhibiting the aggregation of blood platelets. Fibrinogen receptor antagonists of this invention are illustrated by compounds having the formula:



wherein G is




wherein:


A, B, C and D independently represent a carbon atom or a nitrogen atom;

E is $-(\text{CHR}^1)_m-(\text{CHR}^2)_n-\text{F}-(\text{CHR}^3)_o-(\text{CHR}^4)_p$;
 $-(\text{CHR}^1)_m-\text{CR}^2=\text{CR}^3-(\text{CHR}^4)_n-\text{F}$; or
 $-\text{F}-(\text{CHR}^1)_m-\text{CR}^2=\text{CR}^3-(\text{CHR}^4)_n$.

wherein

m, n, o, and p are integers chosen from 0-2; and F is an optional substituent, which when present is chosen from:







5

X is

10



15



or a 4- to 10- membered mono- or polycyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O and S and either unsubstituted or substituted with R¹, R², R³ or R⁴, wherein
 20 R¹, R², R³ and R⁴ are independently selected from the group consisting of hydrogen,


C₁₋₁₀ alkyl,
 aryl C₀₋₈ alkyl,
 oxo,
 thio,
 25 amino C₀₋₈ alkyl, C₁₋₃ acylamino C₀₋₈ alkyl,
 C₁₋₈ alkylamino C₀₋₈ alkyl,
 C₁₋₈ dialkylamino C₀₋₈ alkyl,
 C₁₋₄ alkoxy C₀₋₈ alkyl,
 carboxy C₀₋₈ alkyl, C₁₋₃ alkoxycarbonyl C₀₋₆ alkyl,
 30 carboxy C₀₋₈ alkyloxy and hydroxy C₀₋₈ alkyl;

Y is


C₀₋₈ alkyl,
 C₀₋₈ alkyl-NR³-CO-C₀₋₈ alkyl,
 35 C₀₋₈ alkyl-CONR³-C₀₋₈ alkyl,
 C₀₋₈ alkyl-O-C₀₋₈ alkyl,
 C₀₋₈ alkyl-S(O)_n-C₀₋₈ alkyl, or
 C₀₋₈ alkyl-SO₂-NR³-C₀₋₈ alkyl-,
 C₀₋₈ alkyl-NR³-SO₂-C₀₋₈ alkyl-,
 40 C₁₋₈ alkyl-CO-C₀₋₈ alkyl;

Z is


45



50



55



wherein m is 0-6;

- 5 **R⁵ is** hydrogen
C₁₋₆ alkyl,
C₀₋₆ alkylcarboxy C₀₋₆ alkyl,
C₀₋₆ alkyloxy C₀₋₆ alkyl,
hydroxy C₀₋₆ alkyl,
aryl C₀₋₆ alkyl, or
10 halogen;
- R⁶ is** hydrogen,
C₁₋₈ alkyl,
aryl C₀₋₆ alkyl,
15 C₃₋₈ cycloalkyl C₀₋₆ alkyl,
C₀₋₆ alkylcarboxy C₀₋₆ alkyl, carboxy C₀₋₆ alkyl,
C₁₋₄ alkyloxy C₀₋₆ alkyl,
hydroxy C₀₋₆ alkyl, provided that any of which groups may be substituted or unsubstituted indepen-
20 dently with R¹ or R²,
and provided that, when two R⁶ groups are attached to the same carbon, they may be the same or different;
- R⁷ is** hydrogen, fluorine
C₁₋₈ alkyl,
25 C₃₋₈ cycloalkyl,
aryl C₀₋₆ alkyl,
C₀₋₆ alkylamino C₀₋₆ alkyl,
C₀₋₆ dialkylamino C₀₋₆ alkyl,
C₁₋₈ alkylsulfonylamino C₀₋₆ alkyl,
30 aryl C₀₋₆ alkylsulfonylamino C₀₋₆ alkyl,
C₁₋₈ alkyloxycarbonylamino C₀₋₆ alkyl,
aryl C₀₋₆ alkyloxycarbonylamino C₀₋₆ alkyl,
C₁₋₈ alkylcarbonylamino C₀₋₆ alkyl,
35 aryl C₀₋₆ alkylcarbonylamino C₀₋₆ alkyl,
C₀₋₈ alkylaminocarbonylamino C₀₋₆ alkyl,
aryl C₀₋₆ alkylaminocarbonylamino C₀₋₆ alkyl,
C₁₋₆ alkylsulfonyl C₀₋₆ alkyl,
aryl C₀₋₆ alkylsulfonyl C₀₋₆ alkyl,
40 C₁₋₆ alkylcarbonyl C₀₋₆ alkyl,
aryl C₀₋₆ alkylcarbonyl C₀₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino C₀₋₆ alkyl,
aryl C₀₋₆ alkylthiocarbonylamino C₀₋₆ alkyl
- 45 wherein groups may be unsubstituted or substituted with one or more substituents selected from R¹ and R²,
and provided that when two R⁷ groups are attached to the same carbon atom, they may be the same or dif-
ferent;
- R⁸ is** hydroxy,
C₁₋₈ alkyloxy,
50 aryl C₀₋₆ alkyloxy,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyloxy,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyloxy, or
an L- or D-amino acid joined by an amide linkage and wherein the carboxylic acid moiety of said
amino acid is as the free acid or is esterified by C₁₋₆ alkyl.

55 When substituent R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸ or Y includes the definition CO, (e.g. aryl C₀ alkyl), the group modified by C₀ is not present in the substituent.

"Aryl" means a mono- or polycyclic system composed of 5- and 6- membered aromatic rings containing 0, 1, 2, 3 or 4 heteroatoms chosen from N, O or S and either unsubstituted or substituted with R¹.


"Alkyl" means straight or branched chain alkane, alkene or alkyne.

"Halogen" includes fluorine, chlorine, iodine and bromine.

"Oxo" means =O.

"Thio" means =S.

5 A preferred embodiment of the present invention is



II wherein:

20 E is: $-(CHR^1)_m-(CHR^2)_n-F-(CHR^3)_o-(CHR^4)_p-$,
 $-(CHR^1)_m-CR^2=CR^3-(CHR^4)_n-F-$, or
 $-F-(CHR^1)_m-CR^2=CR^3-(CHR^4)_n-$,
 where m, n, o and p are integers 0-2.


F is an optional substituent which when present is chosen from:

25 0; $-CR^1R^2-$; $-\overset{\overset{O}{\parallel}}{C}-$; $-\overset{\overset{S}{\parallel}}{C}-$; $-\overset{\overset{O}{\parallel}}{C}NR^1$;

30 $-\overset{\overset{O}{\parallel}}{NR^1}C-$; $-\overset{\overset{O}{\parallel}}{C}-O-$; $-O-\overset{\overset{O}{\parallel}}{C}-$;

or $-NR^1R^2$; and X, Y, R^1 , R^2 , R^3 , R^4 , R^6 , R^7 and R^8 are as previously defined.

35 A more preferred embodiment of the present invention is III wherein:



45 E is: $-(CHR^1)_m-F-(CHR^2)_n-$,
 $-CR^1=CR^2-F-$, or
 $-F-CR^1=CR^2-$,
 where m and n are integers 0-2
 and

50 F is an optional substituent which when present is chosen from:

55 0; $-CR^1R^2-$; $-\overset{\overset{O}{\parallel}}{C}-$; $-\overset{\overset{O}{\parallel}}{C}NR^1-$;

$-\overset{\overset{O}{\parallel}}{NR^1}C-$;

or -NR¹R²-

X is -NR¹R² or a 4- to 10-membered mono- or polycyclic aromatic or non-aromatic ring system containing 0, 1 or 2 heteroatoms chosen from N or O and either unsubstituted or substituted with R¹ and R², wherein

5 R¹ and R² are independently chosen from:




hydrogen,
C₁₋₈ alkyl,
aryl C₀₋₆ alkyl,
carboxy C₀₋₆ alkyl,
10 hydroxy C₀₋₆ alkyl,
C₁₋₃ alkyloxy C₀₋₆ alkyl, or
amino C₀₋₆ alkyl;

15 Y is C₀₋₆ alkyl,
C₁₋₆ alkyl-CO-C₀₋₆ alkyl, or
C₀₋₆ alkyl-NR³-CO-C₀₋₆ alkyl;


R⁶ and R⁷ are as previously defined and

20 R⁸ is hydroxy,
C₁₋₈ alkyloxy,
aryl C₁₋₄ alkyloxy, or
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyloxy.


Preferred compounds of the invention are:



5




10



15

20


25



30

35

40




45

50


55

5

10




15




20

25



30

35




40

and


45

50




Generally, compounds of the present invention can be made according to a procedure including the following steps:

a) preparing a triflate activated aromatic group of the following general formula:




10 using




and Ti_2O ;


b) inserting a carbonyl group for the triflate group using metal catalyzed carbonyl insertion, followed by trapping with methanol, to form



c) brominating the heterocyclic methyl group to form




d) cyclizing with a primary amine to form




55 wherein X is an N-terminus protected primary amine, or a primary amine protected directly following this cyclization step;

e) converting the C-terminus ester, via hydrolysis, to an acid



10 f) coupling the acid with an unsubstituted or substituted amino acid or C-terminus protected analog, or diamino acid or C-terminus protected analog, and optionally functionalizing the amino acid at the alpha- or beta-position, to form




and


g) deprotecting the protected C-terminus and N-terminus.

Preferably the procedure involves

25 a) preparing an activated aryl group:




using




and T₂O;


b) inserting a carbonyl group for the triflate group using metal catalyzed carbonyl insertion followed by trapping with methanol to form



c) brominating the aryl methyl group to form




d) cyclizing with a primary amine to form




wherein X is an N-terminus protected primary amine, or a primary amine protected directly following this cyclization step;

25 e) converting the C-terminus ester, via hydrolysis, to an acid



f) coupling the acid with an unsubstituted or substituted amino acid or C-terminus protected analog, or diamino acid or C-terminus protected analog, and optionally functionalizing the amino acid at the alpha- or beta-position via acylation or sulfonylation, to form



and

g) deprotecting the protected C-terminus and N-terminus.





50 An ADP-stimulated platelet aggregation assay was used to determine inhibition associated with compounds of the invention.

Human platelets were isolated from fresh blood, collected into acid citrate/dextrose by differential centrifugation followed by gel filtration on Sepharose 2B in divalent ion-free Tyrode's buffer (pH 7.4) containing 2% bovine serum albumin. Platelet aggregation was measured at 37°C in a Chronolog aggregometer. The reaction mixture contained gel-filtered human platelets (2×10^8 per ml), fibrinogen (100 µg/ml), Ca^{2+} (1 mM), and the compound to be tested. Aggregation was initiated by adding 10 µM ADP 1 minute after the other components had been added. The reaction was allowed to proceed for at least 2 minutes. The extent of inhibition of aggregation was expressed as the percentage of the rate of aggregation observed in the absence of inhibitor.

The IC_{50} is the dose of a particular compound inhibiting aggregation by 50% relative to a control lacking the compound.

Additional preferred embodiments of the invention, shown below with platelet aggregation inhibition potency data (IC_{50} μ m), are shown below.

5





| <u>Compound</u> | <u>IC_{50} μm</u> |
|---|---|
|  | 0.16 |
|  | 4.3 |
|  | 5.7 |
|  | 0.42 |

L-isomer

45

50

55

| Compound | IC ₅₀ μ M |
|---|--------------------------|
|  | 1.0 |
|  | 21 |
| Compound | IC ₅₀ μ M |
|  | 0.92 |
|  | 14.0 |

The abbreviations listed below are defined as Bn, benzyl; NMM, N-methylmorpholine; HOBt, 1-hydroxybenzotriazole; EDC, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; DMF, dimethylformamide; Pib, 4-(4-piperidyl)butanoyl; pTSA, para-toluenesulfonic acid; DMS, dimethylsulfide; TFA, trifluoroacetic acid; THF, tetrahydrofuran; DIBAL, diisobutylaluminum hydride; Boc (or BOC), tert-butoxycarbonyl; Cbz, benzyloxycarbonyl; Suc, succinoyl; alpine borane, β -isopinocampheyl-9-borabicyclo[3.3.1]nonane; TBDMS, tert-butyldimethylsilyl; Jones reagent, chromic acid; NBS, N-Bromosuccinimide; BPO, Benzoyl peroxide; PPh₃, triphenyl phosphine; DMSO, Dimethylsulfoxide; Et₃N, triethylamine; Tf₂O, triflic anhydride; DMAP, 4-dimethylaminopyridine; BOP, benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate; PhCHO, benzaldehyde; and Boc₂O, di-t-butyldicarbonate; dppp, 1,3-bis(diphenylphosphino)propane; ETOH, ethyl acetate; CH₂Cl₂, methylene chloride; HOAc, acetic acid; CH₃OH, methanol; CHCl₃, chloroform.

Unless otherwise indicated, all degree values are Celsius.

The pharmaceutically acceptable salts of the compounds of Formula I include the conventional non-toxic salts or the quarternary ammonium salts of the compounds of Formula I formed, e.g., from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the compounds of Formula I which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts are prepared by reacting the free base or acid with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid or base in a suitable solvent or various combinations of solvents.

5 The pharmaceutically acceptable salts of the acids of Formula I are also readily prepared by conventional procedures such as treating an acid of Formula I with an appropriate amount of a base, such as an alkali or alkaline earth metal hydroxide e.g. sodium, potassium, lithium, calcium, or magnesium, or an organic base such as an amine, e.g., dibenzyl-ethylenediamine, trimethylamine, piperidine, pyrrolidine, benzylamine and the like, or a quaternary ammonium hydroxide such as tetramethylammonium hydroxide and the like.

10 The compounds of Formula I are useful in inhibiting the binding of fibrinogen to blood platelets, inhibiting aggregation of blood platelets, treatment of thrombus formation or embolus formation, and in the prevention of thrombus formation or embolus formation. These compounds are useful as pharmaceutical agents for mammals, especially for humans. The compounds of this invention may be administered to patients where prevention of thrombosis by inhibiting binding of fibrinogen to the platelet membrane glycoprotein complex IIb/IIIa receptor is desired. Compounds of this invention may also be used to prevent or modulate the progress of myocardial infarction, unstable angina and thrombotic stroke, in either acute or chronic settings. In addition, they may be useful in surgery on peripheral arteries (arterial grafts, carotid endarterectomy) and in cardiovascular surgery where manipulation of arteries and organs, and/or the interaction of platelets with artificial surfaces, leads to platelet aggregation and consumption. The aggregated platelets may form thrombi and thromboemboli. Compounds of this invention may be administered to surgical patients to prevent the formation of thrombi and thromboemboli.

Extracorporeal circulation is routinely used for cardiovascular surgery in order to oxygenate blood. Platelets adhere to surfaces of the extracorporeal circuit. Adhesion is dependent on the interaction between GPIIb/IIIa on the platelet membranes and fibrinogen adsorbed to the surface of the circuit. (Glusko et al., 25 *Amer. J. Physiol.*, 1987, 252:H, pp 615-621). Platelets released from artificial surfaces show impaired hemostatic function. Compounds of this invention may be administered to prevent adhesion.

Other applications of these compounds include prevention of platelet thrombosis, thromboembolism, reocclusion, and restenosis during and after thrombolytic therapy and prevention of platelet thrombosis, thromboembolism, reocclusion and restenosis after angioplasty of coronary and other arteries and after coronary artery bypass procedures.

30 The compounds of Formula I may be administered to mammals, preferably in combination with pharmaceutically-acceptable carriers or diluents, optionally with known adjuvants such as alum, in a pharmaceutical composition which is non-toxic and in a therapeutically effective amount, according to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including intravenous, intramuscular, 35 intraperitoneal, trans-dermal, subcutaneous and topical administration.

For oral use of a fibrinogen receptor antagonist according to this invention, the selected compounds may be administered, for example, in the form of tablets or capsules, or as an aqueous solution or suspension. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch, and lubricating agents, such as magnesium stearate, are commonly added. For oral administration in capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or 40 flavoring agents may be added.

For intramuscular, intraperitoneal, subcutaneous, and intravenous use, sterile solutions of the active ingredient are usually prepared, and the pH of the solutions should be suitably adjusted and buffered. For intravenous use, the total concentration of solutes should be controlled in order to render the preparation isotonic.

The present invention also encompasses a pharmaceutical composition useful in the treatment and prevention of diseases related to platelet aggregation, fibrin formation, and thrombus and embolus formation, comprising the administration of a therapeutically effective but non-toxic amount of the compounds of Formula I, with or without pharmaceutically acceptable carriers or diluents.

50 Compositions of this invention include fibrinogen receptor antagonist compounds of this invention in combination with pharmacologically acceptable carriers, e.g. saline, at a pH level e.g. 7.4, suitable for achieving inhibition of platelet aggregation. The compositions may also be combined with anticoagulants such as heparin or warfarin. The compositions may also be combined with thrombolytic agents such as plasminogen activators or streptokinase in order to inhibit platelet aggregation in more acute settings. The composition may further 55 be combined with antiplatelet agents such as aspirin. The compositions are soluble in an aqueous medium, and may therefore be effectively administered in solution.

When a compound according to Formula I is used as a fibrinogen receptor antagonist in a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying

according to the age, weight, and response of the individual patient, as well as the severity of the patients symptoms.

In one exemplary application, a suitable amount of compound is administered orally to a heart attack victim subsequent to angioplasty. Administration occurs subsequent to angioplasty, and is in an amount sufficient to inhibit platelet aggregation, e.g. an amount which achieves a steady state plasma concentration of between about 0.01-50 μM preferably between about 0.01-10 μM .


The present invention also includes a pharmaceutical composition comprising compounds of the present invention in combination with tissue type plasminogen activator or streptokinase. The invention also includes a method for promoting thrombolysis and preventing reocclusion in a patient which comprises administering to the patient an effective amount of compositions of the invention.

The present invention provides a method of inhibiting the binding of fibrinogen to blood platelets, inhibiting aggregation of blood platelets, treating thrombus formation or embolus formation, and in preventing thrombus formation or embolus formation in a mammal, comprising the administration of a therapeutically effective but non-toxic amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents.


The present invention still further provides a method of inhibiting the binding of fibrinogen to blood platelets, inhibiting aggregation of blood platelets, treating thrombus formation or embolus formation, and in preventing thrombus formation or embolus formation in a mammal, comprising the administration of a therapeutically effective but non-toxic amounts of the compounds of this invention in combination with thrombolytic agents, such as tissue plasminogen activators or streptokinase, anticoagulants such as heparin or warfarin, or antiplatelet agents such as aspirin, with or without pharmaceutically acceptable carriers or diluents.

The compounds of Formula I are prepared according to the reaction schemes set forth below.

SCHEME 1




SCHEME 1 CONT'D





55

¹H NMR (300 MHz, CDCl₃) δ 2.18 (3H, s), 3.85 (3H, s), 7.30 (1H, d), 7.84 (1H, s), 7.90 (1H, d).




1 - 3

Dimethyl 4-methylbenzene-1,3-dicarboxylate (1-3)

A solution of 1-2 (30.0 g, 0.121 moles) in methanol/300 ml was treated successively with DMSO (180 ml), triethylamine (0.278 moles), palladium acetate (0.807 g, 3.6 mmoles) and dppp (1.48 g, 3.6 mmoles) as the reaction turned to a clear dark brown solution. Carbon monoxide was then bubbled through the reaction mixture for 3 minutes and the resulting mixture was heated at reflux, while continuing to bubble CO. After refluxing for 4 hours the reaction mixture was concentrated and the resulting brown oil was purified by flask chromatography on silica gel eluting with hexane(90)/EtOAc(10) to provide pure 1-3.

¹H NMR (300 MHz, CDCl₃) δ 2.69 (3H, s), 3.95 (3H, s), 3.96 (3H, s), 7.37 (1H, d), 8.09 (1H, dd), 8.60 (1H, d).




1 - 4

Dimethyl 4-bromomethylbenzene-1,3-dicarboxylic acid (1-4)

A solution of 1-3 (1.35 g, 6.5 mmole) in CHCl₃ (20 ml) was treated with dibenzoyl peroxide (0.078g, 3.5 mmol) and N-bromosuccinimide (NBS) (1.1g, 6.5 mmole) and the resulting solution was heated at reflux for 2 hours.

The cooled reaction mixture was concentrated, taken up in CCl₄, filtered and the filtrate was concentrated to give 1-4 as a tan solid. R_f 0.5 [silica gel, hexane(70)/EtOAc(30)].

Preparation of Boc-4-Piperidine-2-ethanol (1-5)



1 - 5


4-Piperidine-2-ethanol (Aldrich) (130 g, 1.0 mole) was dissolved in 700 mL dioxane, cooled to 0°C and treated with 3 N NaOH (336 mL, 1.0 mole), and di-*t*-butyldicarbonate (221.8 g, 1.0 mole). The ice bath was removed and the reaction stirred overnight. The reaction was concentrated, diluted with water and extracted with ether. The ether layers were combined, washed with brine, dried over MgSO₄, filtered and evaporated to give 1-5.

R_f = 0.37 in 1:1 EtOAc/Hexanes, ninhydrin stain


¹H NMR (300MHz, CDCl₃) δ 4.07 (bs, 2H), 3.7 (bs, 2H), 2.7 (t, J = 12.5 Hz, 2H), 1.8-1.6 (m, 6H), 1.51 (s, 9H), 1.1 (ddd, J = 4.3, 12.5, 12 Hz, 2H).

Boc-4-piperidine-2-ethyl iodide (1-6)


Boc-4-piperidine-2-ethanol (1-5) (10.42 g, 0.048 mole) was dissolved in 400 ml benzene and imidazole (4.66 g, 0.068 moles) and triphenyl-phosphine (15.24 g, 0.05 moles) were added at room temperature. After 6 hours the reaction mixture was filtered and the filtrate was evaporated to give a dark residue. This was purified by flash chromatography on silica gel eluting with 10% EtOAc-hexanes to give 1-6 as a yellow oil.

Boc-4-piperidine-2-ethylazide (1-7)

To 1-6 (27.9 g, 0.082 moles) dissolved in DMSO (400 ml) was added sodium azide (5.01 g, 0.086 moles) at room temperature and the resulting solution was heated at 65° for 2 hours. The cooled reaction mixture was diluted with 250 ml EtOAc, extracted with 2 x 100 ml portions of water 2 x 50 ml portions of brine and then dried (MgSO₄). Solvent removal provided 1-7 as a pale yellow oil, R_f 0.5 (silica gel, 70% acetone/hexane).

Boc-4-piperidine-2-ethylamine(1-8)

To a solution of 1-5 (19.3 g, 0.076 moles) in THF (400 ml)/H₂O (195 ml) was added triphenyl-phosphine (80.0g, 0.305 moles) in one portion at room temperature. This was stirred at room temperature 3 hours and the organic solvents were then removed *in vacuo*. The residue was acidified to pH 2 with 10% KHSO₄ solution and this was extracted 4 x 100 ml portions of EtOAc. The organic extract was extracted with 2 x 100 ml portions of 10% KHSO₄ and the aqueous phases were combined and the pH was adjusted to 10 with 2N NaOH. This solution was extracted with 4 x 200 ml portions of CH₂Cl₂. These were combined, dried (MgSO₄) and the solvent was removed to give 1-8 as an oil. R_f 0.3 (silica gel, eluting with 10% CH₃OH in CHCl₃/NH₃). ¹H NMR (300 MHz, CDCl₃) δ 4.05 (broad, 2H), 2.72 (t, J=7.2Hz, 2H), 2.62 (m, 2H), 1.64 (d, J=12.2Hz, 2H), 1.43 (s, 9H), 1.42-1.32 (m, 5H), 1.09 (m, 2H).




Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (1-9)

A solution of 1-4 (1.0 g, 3.5 mmoles) in benzene (5 ml) was treated with 1-8 (0.80 g, 3.5 mmol) and triethylamine (0.49 ml, 3.5 mmol) and the reaction mixture was heated at reflux for 3 hours. The solvent was removed and the residue was taken up in EtOAc, washed in 10% KHSO₄ solution, H₂O, brine and dried. Solvent removal gave a residue that was purified by flash chromatography on silica gel eluting with hexane(1)/EtOAc(1) to give pure 1-9.


R_f 0.2 (silica gel, hexane(1)/EtOAc(1)).

¹H NMR (300 MHz, CDCl₃) δ 1.08 (2H, m), 1.43 (9H, s), 1.61 (4H, m), 1.73 (2H, bd), 2.62 (2H, bt), 3.64 (2H, t), 3.93 (3H, s), 4.07 (2H, m), 4.40 (2H, s), 7.50 (1H, d), 8.21 (1H, dd), 8.47 (1H, d).

1 - 1 01-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (1-10)

A solution of 1-9 (0.43 g, 1.12 mmole) in THF (1)/MeOH(1)/H₂O(1) (9 ml) was treated at room temperature with LiOH·H₂O (0.235 g, 5.6 mmol) and the resulting solution was stirred for 4 hours. The reaction mixture was then diluted with EtOAc (75 ml)/10% KHSO₄ solution (30 ml) and the organic phase was separated and dried (Na₂SO₄). Solvent removal gave the desired acid 1-10. R_f 0.5 (silica gel, CH₂Cl₂(9)/MeOH (0.5)/HOAc(0.5)).


¹H NMR (300 MHz, CDCl₃) δ 1.12 (2H, m), 1.42 (9H, s), 1.60 (3H, m), 1.71 (2H, bd), 2.63 (2H, bt), 3.68 (2H, t), 4.08 (2H, m), 4.40 (2H, s), 7.03 (1H, d), 8.28 (1H, dd), 8.60 (1H, s).

1 - 1 11-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(carboethoxy)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (1-11)

A solution of 1-10 (0.35 g, 0.94 mmole), triethylamine (0.40 ml, 2.82 mmol), and β-alanine ethyl ester (0.22 g, 1.41 mmol) (Aldrich) in CH₃CN (5 ml) was treated at room temperature with BOP (1.2 mmoles) reagent and the resulting solution was stirred for 16 hours.

The solvent was removed and the residue was taken up in EtOAc, washed with H₂O, 10% KHSO₄ solution, brine and dried (Na₂SO₄). Solvent removal gave a residue that was purified by flash chromatography on silica gel eluting with hexane(20)/EtOAc(80) to give pure 1-11 as a clear oil.



¹H NMR (300 MHz, CDCl₃) δ 1.10-1.30 (3H, m), 1.44 (9H, s), 1.60 (3H, m), 1.75 (2H, bd), 2.63 (4H, m), 3.70 (4H, m), 4.05-4.20 (4H, m), 4.38 (2H, s), 7.50 (1H, d), 8.08 (2H, m).

1 - 1 21-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxy-ethyl)-2-[2-(4-piperidiny)ethyl]-3-oxo (1-12)

A solution of 1-11 (0.32 g, 0.68 mmol) in THF(1)/MeOH(1)/H₂O(1) (9 ml) was treated with LiOH·H₂O (0.14 g, 3.4 mmoles) at room temperature for 1.0 hr. The solvent was then removed and the residue was taken up in EtOAc and washed with 10% KHSO₄ solution, brine and dried (Na₂SO₄). Solvent removal gave the desired acid. R_f 0.3 (silica gel, CHCl₃ (9)/MeOH (0.5)/HOAc (0.5)).

This acid (0.30 g, 0.68 mmole) was dissolved in CH₂Cl₂ and anisole (150 μl) was added. This was cooled to -15°C and trifluoroacetic acid (3 ml) was added and the resulting mix stirred for 0.5 hours. The solvent was removed and the residue purified by flash chromatography on silica gel eluting with EtOH (9)/NH₄OH (1.2)/H₂O (1.2) to provide pure 1-12.


¹H NMR (300 MHz, D₂O) δ 1.30 (7H, m), 1.50-1.70 (3H, m), 1.83 (2H, bd), 2.38 (2H, t), 2.80 (2H, dt), 3.27 (2H, bd), 3.50 (4H, m), 4.42 (2H, s), 7.51 (1H, d), 7.83 (2H, m).

1 - 1 11 - 1 31-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(carboethoxy)ethyl]-2-[2-(4-piperidiny)ethyl]-3-oxo (1-13)

A solution of 1-11 (0.72 g, 1.57 mmoles) in EtOAc (20 ml) was cooled to -78°C and HCl gas was bubbled through. This solution for 1-2 minutes and the reaction mixture was then stirred at 0°C. After a few minutes a white solid had precipitated and this mixture was stirred for 0.5 hours. The solvent was then removed and the residue was triturated with Et₂O to give pure 1-13.

¹H NMR (300 MHz, CD₃OD) δ 1.23 (3H, t), 1.45 (2H, m), 1.66 (2H, m), 1.72 (2H, m), 2.07 (2H, m), 2.65 (2H, t), 2.94 (2H, m), 3.47 (2H, bd), 3.68 (4H, m), 4.12 (2H, q), 4.57 (2H, s), 7.67 (1H, d), 8.03 (1H, dd), 8.14 (1H,

d).




1H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butylcarbonyloxymethylcarboxy)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (1-15)

35 A slurry of 1-16 (0.80 g, 1.8 mmoles) in MeOH (20 ml) was treated with Cs_2CO_3 (0.24 g, 0.90 mmoles) at room temperature and the resulting mixture was stirred for 45 minutes. The solvent was then removed and the residue was slurried in DMF (20 ml) and this was treated at room temperature with chloromethyl pivalate (1.8 mmoles). The resulting mixture was stirred at room temperature for 24 hours.

40 The reaction mixture was then diluted with EtOAc and washed with H_2O , 10% KHSO_4 , saturated with NaHCO_3 solvent and brine. The organic phase was dried (MgSO_4), and the solvent removed to provide 1-15 as a white solid.

^1H NMR (300 MHz, CDCl_3) δ 1.11-1.25 (13H, m), 1.46 (9H, s), 1.63 (2H, q), 1.77 (2H, bd), 2.62-2.76 (4H, m), 3.72 (9H, m), 4.09 (2H, bd), 4.42 (2H, s), 5.80 (2H, s), 6.89 (1H, bt), 7.53 (1H, d), 8.09 (1H, d), 8.14 (1H, s).




55 1-H-Isoindole-5-carbox-amide, 2,3-dihydro-N-[2-(t-butylcarbonyloxymethylcarboxy) ethyl]-2-[2-(4-piperidinyl) ethyl]-3-oxo (1-16)


A solution of 1-15 (15 mg) in EtOAc (5 ml) was cooled to -78°C and treated with HCl gas for 10 minutes

and the resulting solution was stirred at -10°C for 1.0 hour. The solvent was then removed to provide pure 1-16 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.06 (9H, s), 1.92 (1H, m), 1.70 (2H, m), 2.08 (2H, bd), 3.73 (2H, t), 2.95 (2H, dt), 3.38 (2H, bd), 3.70 (6H, m), 4.58 (2H, s), 5.86 (2H, s), 7.67 (1H, d), 8.06 (1H, d), 8.17 (1H, s).




1-14




1-17




LiOH



1-18



HCl (gas)
EtOAc




1-19

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[L-Phe(OEt)-2-(carboxamido)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo(1-17)

1-14 (0.35 g, 0.76 mmol) was treated with L-phenylalanine ethyl ester (2.0 mmol), N-methylmorpholine (2.0 mmol) and BOP (0.886 g, 2.0 mmol), in CH₃CN (5 ml) at room temp for 24 hrs. as described for 6-3. Flash chromatography on silica gel eluting with EtOAc (9)/MeOH (1) gave pure 1-17 as a white solid. R_f 0.3 (silica gel, CHCl₃(2)/acetone (1).

¹H NMR (300 MHz, CDCl₃) δ 1.28 (3H, t), 1.47 (9H, s), 1.79 (2H, bd), 2.54 (2H, t), 2.72 (2H, m), 3.15 (2H, m), 3.75 (5H, m), 4.20 (4H, m), 4.43 (2H, s), 2.90 (1H, q), 7.12 (2H, m), 7.25 (5H, m), 7.54 (1H, d), 8.08 (1H, d), 8.19 (1H, s).




1-18

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[L-Phe-2-(carboxamido)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo(1-18)

1-17 (0.46 g, 0.72 mmol) was treated with LiOH·H₂O (0.152 g, 3.6 mmol) as described for 1-12 to give 1-18 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.13 (2H, m), 1.43 (9H, s), 1.66 (2H, q), 1.80 (2H, bd), 2.50 (2H, t), 2.70 (2H, m), 2.93 (1H, m), 3.20 (1H, dd), 3.58 (2H, q), 3.70 (2H, t), 4.04 (2H, m), 4.56 (2H, s), 4.68 (1H, m), 7.20 (5H, m), 7.56 (1H, d), 8.02 (1H, d), 8.15 (1H, s).




1-19



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[L-Phe-2-(carboxamido)ethyl]-2-[2-(4-piperidinyl)ethyl]-3-oxo(1-19)

1-18 (0.35 g, 0.37 mmol) was treated with HCl gas as described for 1-13 to give pure 1-19 as a white solid.

¹H NMR (300 MHz, D₂O) δ 1.35 (2H, m), 1.62 (2H, m), 1.93 (2H, m), 2.43 (2H, m), 2.79 (3H, m), 3.07 (1H, m), 3.28 (2H, m), 3.45 (2H, m), 4.50 (2H, s), 6.80 (1H, m), 6.92 (2H, m), 7.00 (2H, m), 7.55 (1H, d), 7.77 (2H, bs).




1-14



1-20

1. LiOH·H₂O
2. HCl (gas)/EtOAc




1-21

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[L-Pro(OC₂H₅)-2-(carboxamido)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (1-20)

1-14 (0.35 g, 0.76 mmoles) was treated with L-Proline ethyl ester (0.288 g, 2.0 mmoles), N-methylmorpholine (2.0 mmoles) and BOP (0.886 g, 2.0 mmoles) in CH₃CN (5 ml) as described for 1-17 to give an oily residue. This was purified by flash chromatography on silica gel eluting with acetone (1)/CHCl₃(1) to give pure 1-20.

¹H NMR (300 MHz, CDCl₃) δ 1.16 (2H, m), 1.45 (9H, s), 1.42 (2H, q), 1.65 (2H, bd), 2.03 (2H, m), 2.66 (5H, m), 3.51 (1H, m), 3.67 (2H, m), 3.80 (2H, m), 4.09 (2H, m), 4.20 (2H, q), 4.40 (2H, s), 4.50 (1H, m), 7.41 (1H, m), 7.50 (1H, d), 8.03 (1H, d), 8.19 (1H, s).



1-21

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[L-Pro-2-(carboxamido)ethyl]-2-[2-(4-piperidinyl)ethyl]-3-

oxo (1-21)


1-20 (0.2 g, 0.34 mmoles) was treated with LiOH·H₂O (0.071 g, 1.7 mmoles) as described for 1-12 to give the desired acid.


¹H NMR (300 MHz, CD₃OD) δ 1.15 (2H, m), 1.44 (9H, s), 1.67 (2H, q), 2.80 (2H, bd), 2.25 (1H, m) 2.73 (2H, m), 3.68 (4H, m), 4.06 (2H, m), 4.55 (2H, s), 7.66 (1H, d), 8.05 (1H, d), 8.17 (1H, s).

This acid (0.15 g) was dissolved in EtOAc (10 ml) and treated with HCl gas as described for 1-13 to give pure 1-21 as a white solid.

¹H NMR (300 MHz, D₂O) δ 1.48 (2H, m), 1.67 (1H, m), 1.76 (2H, m), 2.06 (4H, m), 2.32 (1H, m), 2.62 (1H, m), 2.84 (2H, t), 2.96 (2H, t), 3.43 (2H, d), 3.70 (6H, m), 4.47 (1H, m), 4.66 (2H, s), 7.72 (1H, d), 8.00 (1H, d), 8.09 (1H, s).

SCHEME 2






4-(N-t-Butyloxycarbonylpiperidinyl)methylamine (2-3)

55 A solution of 4-(piperidinyl)methylamine (2-1) (22.8 g, 0.2 mmoles) in toluene (250 ml) was treated with benzaldehyde (21.2 g, 0.2 mmoles) at room temperature and the resulting mixture was heated at reflux for 3 hours with the aid of a Dean-Stark trap for water removal. The cooled reaction mixture containing the desired Schiff's base 2-2 was treated portionwise with di-t-butyl dicarbonate (47.96 g, 0.22 moles) and the resulting

solution was stirred at room temperature for 16 hours. The solvent was then removed and the residue was cooled to 0-5°C and treated with 1N KHSO₄ (220 ml) with stirring for 3 hours. The resulting reaction mixture was extracted with ether (3 x 200 ml) and then made basic with 1N KOH solution and extracted with CHCl₃ (4 x 75 ml). The combined organic extract was washed with brine, dried (Na₂SO₄) filtered through celite, and the solvent removed to provide pure 2-3 as a clear oil.

¹H NMR (300 MHz, CDCl₃) δ 1.13 (2H, m), 1.45 (9H, s), 1.60 (1H, m), 1.74 (2H, d), 2.68 (4H, m), 4.15 (2H, bd).




2-4

Methyl-1H-Isoindole-4-carboxylate, 2,3-dihydro-N-[(4-N-t-butyloxycarbonyl)piperidinyl]methyl]-3-oxo (2-4)

A solution of 1-4 (3.01 g, 10.5 mmoles) in benzene (20 ml) was treated at room temperature with 2-3 (2.30 g, 10.7 mmoles) and Et₃N (10.8 mmoles) and the resulting solution was heated at reflux for 2 hours. The solvent was removed and the residue was taken up in EtOAc (200 ml) and extracted with 10% KHSO₄ solution (5 x 50 ml), brine and dried (MgSO₄). Solvent removal gave a residue that was purified by flash chromatography on silica gel eluting with hexane (1)/EtOAc (1) to give pure 2-4, R_f 0.25.

¹H NMR (300 MHz, CDCl₃) δ 1.29 (2H, m), 1.45 (9H, s), 1.67 (4H, m), 1.95 (1H, m), 2.70 (2H, t), 3.52 (2H, b), 3.97 (3H, s), 4.13 (2H, b), 4.95 (2H, s), 7.52 (1H, d), 8.23 (1H, d), 8.50 (1H, s).



2-5


1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(carboethoxyethyl)-2-[(4-N-t-butyloxycarbonyl)piperidinyl]methyl]-3-oxo (2-5)

A solution of 2-4 (1.92g, 5.58 mmoles) in 150 ml of THF(1)/MeOH(1)/H₂O(1) was treated with LiOH·H₂O (1.20 g, 28.6 mmoles) at room temperature and the resulting solution was stirred for 1.0 hr. The solvent was then removed and the residue was taken up in H₂O (100 ml) acidified to pH 2 with 10% KHSO₄ solution. The desired acid precipitated from solution and was collected.

¹H NMR (300 MHz, CD₃OD) δ 1.13 (2H, m), 1.40 (9H, s), 1.50-1.65 (3H, m), 2.70 (2H, b), 3.45 (2H, d), 3.98 (2H, d), 4.45 (2H, s), 7.60 (1H, d), 8.10 (1H, d), 8.21 (1H, s).

This acid (1.62 g, 4.91 mmoles) was dissolved in CH₃CN (25 ml) and treated at 0° successively with Et₃N (34.4 mmoles), β-alanine ethyl ester (5.0 mmoles), and BOP (3.27 g, 7.38 mmoles). The reaction mixture was then stirred at room temperature for 16 hrs. The solvent was removed and the residue purified by flash chromatography in silica gel eluting with EtOAc (7)/hexane (1) to provide 2-5 as a white solid.

¹H NMR (300 MHz, CDCl₃) δ 1.27 (6H, m), 1.42 (9H, s), 1.67 (5H, m), 1.95 (1H, m), 2.66 (4H, m), 3.50 (2H, b), 3.74 (2H, g), 4.16 (4H, m), 4.45 (2H, s), 7.00 (1H, t), 7.53 (1H, d), 8.11 (2H, m).

2-6

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[(4-piperidiny)methyl]-3-oxo (2-6).


A solution of 2-5 (0.86 g, 2.0 mmoles) in 60 ml of THF(1)/MeOH(1)/H₂O(1) was treated with LiOH·H₂O (0.45 g, 10.7 mmoles) at room temperature and the resulting solution was stirred at room temperature for 1.0 hr. The solvent was removed and the residue was dissolved in H₂O (25 ml), acidified to pH 2-3 with 10% KHSO₄ solution and extracted with EtOAc (4 x 25 ml). The combined organic extracts were washed with brine, dried (Na₂SO₄) and the solvent removed to give the desired acid as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.16 (2H, m), 1.39 (9H, s), 1.45 (1H, m), 1.80 (2H, bd), 1.93 (2H, d), 2.58 (2H, t), 2.70 (2H, b), 3.45 (2H, d), 3.57 (2H, t), 4.00 (2H, m), 7.59 (1H, d), 8.00 (1H, d), 8.09 (1H, s).

This acid (0.80 g, 1.89 mmoles) was treated with HCl gas in EtOAc solution as described for 2-3 to provide pure 2-6 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.43 (2H, m), 1.85 (2H, m), 2.10 (1H, m), 2.56 (2H, t), 2.90 (2H, t), 3.34 (2H, bd), 3.54 (4H, m), 4.52 (2H, s), 7.61 (1H, d), 8.00 (1H, d), 8.10 (1H, s).

2-5 can also be converted to 2-7 as shown below:




1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[(2-carboethoxy)ethyl]-2-[2-(4-piperidiny)ethyl]-3-oxo (2-7).

Treatment of 2-5 (0.90g, 2.09 mmoles) in EtOAc with HCl gas as described for 1-12 gave 2-7 as a white, solid.

¹H NMR (300 MHz, CD₃OD) δ 1.09 (3H, t), 1.45 (2H, m), 1.86 (2H, bd), 2.13 (2H, m), 2.60 (2H, t), 2.90 (2H, t), 3.32 (2H, bd), 3.56 (4H, m), 4.08 (2H, q), 4.56 (2H, s), 7.62 (1H, d), 8.00 (1H, d), 8.09 (1H, s).


SCHEME 3



Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[3-aminopropyl]-3-oxo (3-1)

A solution of 1-4 (2.58 g, 8.99 mmoles in benzene (10 ml) was treated with Et₃N (12.9 mmoles) and 1,3-diaminopropane (13.0 mmoles) at room temperature and the resulting mixture was heated at reflux for 2 hrs. The reaction mixture was cooled and the solvent removed to give 3-1.

¹H NMR (300 MHz, CD₃OD) δ 1.53 (9H, s), 1.79 (2H, m), 3.02 (2H, m), 3.58 (2H, m), 3.84 (3H, s), 4.48 (2H, s), 7.58 (1H, d), 8.10 (1H, d), 8.20 (1H, s).

3-2


1-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[3-(N-t-butyloxycarbonylamino)propyl]-3-oxo (3-2)

3-1 (2.22 g, 8.99 mmoles) was suspended in 100 ml of THF(1)/H₂O(1) and treated with Et₃N (9.3 mmoles) and di-*t*-butyl dicarbonate (4.0 g, 18.3 mmoles) and the resulting mixture was stirred vigorously for 5 hrs. The solvent was removed and the residue was purified by flash chromatography to give the desired protected ester.

¹H NMR (300 MHz, CD₃OD) δ 1.53 (9H, s), 1.80 (2H, m), 3.03 (2H, m), 3.58 (2H, m), 3.86 (3H, s), 4.48 (2H, s), 7.55 (1H, d), 8.10 (1H, d), 8.20 (1H, s).

This ester (0.67 g, 1.93 mmoles) was treated with LiOH·H₂O (0.41 g, 9.76 mmoles) in 60 ml of THF(1)/MeOH(1)/H₂O(1) at room temperature for 1 hr. Solvent removal gave a residue that was dissolved in 25 ml H₂O, acidified to pH 2-3 with 10% KHSO₄ solution and extracted with EtOAc (4x25 ml). The organic extract was washed with brine, dried (MgSO₄) and the solvent removed to give 3-2 as a white solid.


¹H NMR (300 MHz, CD₃OD) δ 1.35 (9H, s), 1.80 (2H, m), 3.04 (2H, t), 3.62 (2H, t), 4.55 (2H, s), 7.62 (1H, d), 8.20 (1H, d), 8.32 (1H, s).

3-3

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl)ethyl]-2-[3-(N-t-butyloxycarbonylamino)propyl]-3-oxo (3-3).

A solution of 3-2 (0.65 g, 1.94 mmoles) in 10 ml CH₃CN was cooled to 0-10° and treated with Et₃N (13.6 mmoles) and BOP (1.30 g, 2.93 mmoles) and the resulting solution was stirred at room temperature for 16 hrs. The solvent was then removed and the residue was taken up in EtOAc (100 ml) extracted with H₂O (4x25 ml), 10% KHSO₄ solution and dried (MgSO₄). Solvent removal gave a residue that was purified by flash chromatography on silica gel eluting with CHCl₃(95)/MeOH(5) to give pure 3-3 as a white solid. R_f 0.3 (silica gel, CHCl₃(95)/MeOH(5)).

¹H NMR (300 MHz, CDCl₃) δ 1.46 (9H, s), 1.53 (9H, s), 1.90 (2H, m), 2.62 (2H, t), 3.60 (2H, m), 3.76 (4H, m), 4.50 (2H, s), 7.00 (1H, t), 7.62 (1H, d), 8.17 (1H, d), 8.20 (1H, s).


3-4


1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[3-aminopropyl]-3-oxo (3-4)

3-3 (0.77g, 1.67 mmoles) was suspended in EtOAc (25 ml) and after cooling to -70° , HCl gas was bubbled into the mixture for 5 minutes at which time the reaction mixture was homogeneous. The reaction mixture was then stirred at $0-5^{\circ}$ for 30 minutes. The solvent was removed and the residue was dried at high vacuum to provide pure 3-4 as a white solid.

^1H NMR (300 MHz, CD_3OD) δ 2.00 (2H, m), 2.60 (2H, t), 2.92 (2H, t), 3.59 (2H, m), 3.70 (2H, t), 4.28 (2H, s), 7.63 (1H, d), 8.02 (1H, d), 8.12 (1H, s).


SCHEME 4




4-1Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[5-aminopentyl]-3-oxo (4-1)

A solution of 1-4 (2.56 g, 8.92 mmoles) in benzene (15 ml) was treated with Et₃N (11.5 mmoles) and 1,5-diaminopentane (11.9 mmoles) and the resulting reaction mixture was heated at reflux for 3 hrs. The solvent was then removed and the residue was purified by flash chromatography on silica gel eluting with 25% MeOH in CHCl₃ (MHz) to provide pure 4-1.

¹H NMR (300 MHz, CDCl₃) δ 1.77 (6H, m), 2.45 (2H, bs), 2.71 (2H, t), 3.63 (2H, t), 4.44 (2H, s), 7.52 (1H, d), 8.22 (1H, d), 8.49 (1H, s).

4-2Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[5-(N-t-butyloxycarbonylamino)pentyl]-3-oxo (4-2)

A solution of 4-1 (0.64 g, 2.32 mmoles) in CH₂Cl₂ (10 ml) was treated at room temperature with Et₃N (2.29 mmoles) and Boc₂O (0.74 g, 3.39 mmoles) for 48 hrs. The solvent was then removed and the residue was purified by flash chromatography on silica gel eluting with hexane(7)/acetone(3) to give pure 4-2.

4-31-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(2-t-butyloxycarbonyl)ethyl]-2-[5-N-t-butyloxycarbonylamino]pentyl]-3-oxo (4-3)


A solution of 4-2 (0.71g, 1.89 mmoles) in THF(1)/MeOH(1)/H₂O(1) (60 ml) was treated with LiOH·H₂O (0.42 g, 10.0 mmoles) at room temperature for 0.5 hr. The solvent was then removed and the residue was dissolved in H₂O (50 ml), acidified to pH 2-3 with 10% KHSO₄ solution and extracted with EtOAc. The organic phase was washed with brine, dried (MgSO₄) and the solvent removed to give the desired acid.

¹H NMR (300 MHz, CD₃OD) δ 1.30 (9H, s), 1.45 (3H, m), 1.63 (3H, m), 2.92 (2H, t), 3.55 (2H, t), 4.47 (2H, s), 7.58 (1H, d), 8.16 (1H, d), 8.03 (1H, s).

This acid (0.75g, 2.07 mmoles) was dissolved in CH₃CN (15 ml) and was treated at room temperature with β-alanine t-butyl ester (0.39g, 2.54 mmoles), BOP (1.4 g, 3.16 mmoles), Et₃N (6.1 mmoles) and the resulting solution was stirred at room temperature for 20 hrs. The solvent was then removed and the residue was dis-

solved in EtOAc and extracted with H₂O, 10% KHSO₄ solution and brine. The organic phase was dried (MgSO₄) and was solvent was removed to give a residue that was purified by flash chromatography on silica gel eluting with EtOAc(7)/hexane(3) to give pure 4-3.

¹H NMR (300 MHz, CD₃OD) δ 1.39 (9H, s), 1.45 (2H, m), 1.65 (2H, m), 2.50 (2H, t), 2.96 (2H, q), 3.53 (4H, q), 4.47 (2H, s), 7.58 (1H, d), 7.96 (1H, d), 8.08 (1H, s).




4-4

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[5-aminopentyl]-3-oxo (4-4)




A solution of 4-3 (0.71g, 1.45 mmoles) in EtOAc (20 ml) was cooled to -78° and treated with HCl gas for 10 minutes. The resulting solution was stirred in at 0° for 0.5 hr. The solvent was removed to provide 4-4 as white solid.

¹H NMR (300 MHz, D₂O) δ 1.29 (2H, m), 1.63 (4H, m), 2.62 (2H, t), 2.87 (2H, t), 3.52 (4H, m), 4.40 (2H, s), 7.51 (1H, d), 7.80 (2H, m).

SCHEME 5



SCHEME 5 cont'd



5-3


N-t-Butyloxycarbonyl-N-methyl-1,3-diaminopropane (5-3).

A solution of N-methyl-1,3-diaminopropane (2.05 g, 23.2 mmoles) in toluene (30 ml) was treated with benzaldehyde (2.41 g, 22.7 mmoles) and the resulting mixture was heated at reflux with use of a Dean-Stark trap. After 2 hrs. the reaction mixture was cooled and treated with Boc₂O (5.57 g, 25.5 mmoles) portionwise and

the resulting solution was stirred for 48 hrs.

The solvent was then removed and the residue was cooled to 0-5° and acidified to pH 2-3 with 10% KHSO₄ solution (25 ml) and the resulting slurry was stirred for 3 hrs. This mixture was then extracted with EtOAc and the aqueous phase was adjusted to pH 9 with 1N NaOH and extracted with CHCl₃ (5x25 ml). The dried organic phase was concentrated to give 5-3 as an oil.

¹H NMR (300 MHz, CDCl₃) δ 1.47 (9H, s), 1.72 (2H, bt), 2.16 (2H, bs), 2.75 (2H, t), 2.87 (3H, s), 3.34 (2H, bs).




5-4

Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[2-(3-N-t-butyloxycarbonyl-N-methylamino)propyl]-3-oxo
(5-4)

A solution of 1-4 (2.0 g, 6.97 mmoles) in benzene (10 ml) was treated with 5-3 (1.19 g, 6.32 mmoles) and Et₃N (7.17 mmoles) and the resulting solution was heated at reflux for 24 hrs. The cooled reaction mixture was then dissolved in EtOAc (150 ml), washed with 10% KHSO₄ solution (4x50 ml), brine (50 ml) and dried (MgSO₄). The solvent was removed to give an oil that was purified by flash chromatography on silica gel eluting with EtOAc(7)/hexane(1) to give pure 5-4 as a white solid.

¹H NMR (300 MHz, CDCl₃) δ 1.45 (9H, s), 1.92 (2H, m), 2.90 (3H, s), 3.30 (2H, t), 3.68 (2H, t), 3.97 (3H, s), 4.50 (2H, s), 7.55 (1H, d), 8.26 (1H, d), 8.52 (1H, s).



5-5


1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl) ethyl]-2-[3-(N-t-butyloxycarbonyl-N-methylamino) propyl]-3-oxo (5-5)

A solution of 5-4 (1.28 g, 3.53 mmoles) in THF(1)/MeOH(1)/H₂(1) (105 ml) was treated with LiOH·H₂O (0.76 g, 18.1 mmoles) and the resulting solution was stirred at room temperature for 30 minutes. The solvent was then removed and the residue was taken up in H₂O (30 ml), acidified to pH 2-3 with 10% KHSO₄ solution, and extracted with EtOAc. The combined organic extracts were washed with brine, dried (MgSO₄) and the solvent removed to provide the desired acid.

¹H NMR (300 MHz, CD₃OD) δ 1.34 (9H, s), 1.86 (2H, m), 2.78 (3H, s), 3.22 (2H, m), 3.55 (2H, t), 4.50 (2H, s), 7.60 (1H, d), 8.17 (1H, d), 8.30 (1H, s).


This acid (1.28 g, 3.59 mmoles) was dissolved in CH₃CN (20 ml) and treated successively with β-alanine t-butyl ester hydrochloride (0.65 g, 3.59 mmoles), Et₃N (2.51 mmoles), and BOP (2.39 g, 5.40 mmoles) and the resulting cloudy suspension was stirred at room temperature for 20 hrs. The reaction mixture was then concentrated and the residue was taken up in EtOAc (100 ml), extracted with H₂O (2x25 ml), 10% KHSO₄ solution (4x25 ml), brine and dried (MgSO₄). Solvent removal gave a residue that was purified by flash chromatography on silica gel eluting with acetone(3)/hexane(7) to give pure 5-5 as a white solid.


¹H NMR (300 MHz, CDCl₃) δ 1.42 (9H, s), 1.44 (9H, s), 1.93 (2H, m), 2.37 (2H, t), 2.88 (3H, s), 3.30 (2H, t), 3.68 (4H, m), 4.47 (2H, s), 6.98 (1H, bt), 7.55 (1H, d), 8.09 (1H, d), 8.12 (1H, s).

5-61-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[3-(N-methylamino)propyl]-3-oxo (5-6)

A solution of 5-5 (1.42 g, 2.09 mmoles) in EtOAc (40 ml) was cooled to -78° and treated with HCl gas for 3-5 minutes. The resulting solution was stirred at 0° for 0.5 hr. The solvent was then removed to provide 5-6 as a white solid.


^1H NMR (300 MHz, D_2O) δ 2.00 (2H, m), 2.62 (5H, m), 3.00 (2H, t), 3.60 (4H, m), 4.29 (2H, s), 7.75 (1H, d), 7.83 (1H, d), 7.88 (1H, s).

SCHEME 6

6-1


Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[6-aminoheptyl]-3-oxo (6-1)

Treatment of 1-4 with 1,6-diaminohexane as described for 1-9 provided 6-1 as a white solid. R_f 0.5 (silica gel, hexane (9)/EtOAc (1)).

6-2


1-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[6-N (t-butyloxycarbonylamino)heptyl]-3-oxo (6-2)

Treatment of 6-1 with Boc_2O (1 equiv) and triethylamine (2 equivalents) in H_2O (1)/THF(1) (100 ml) at room temperature for 48 hours followed by solvent removal gave crude BOC-protected derivative. Hydrolysis of this with $\text{LiOH}\cdot\text{H}_2\text{O}$ (4 equiv.) as described for 1-10 gave 6-2 as an oil. ^1H NMR/(300 MHz, CD_3OD) δ 1.32 (17H, m), 1.68 (2H, m) 2.95 (2H, t), 4.50 (2H, s), 7.62 (1H, d), 8.19 (1H, d), 8.31 (1H, s).

6-3

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl)ethyl]-2-[6-N-(t-butyloxycarbonylamino)heptyl]-3-oxo (6-3)


Treatment of 6-2 (1.18 g, 3.12 mmoles) with t-butyl β -alanine (0.54 g, 3.51 mmoles) as described for 1-11 gave crude 6-3. This was purified by flash chromatography on silica gel eluting with pet ether (6)/EtOAc (4) to provide 6-3 as an oil. R_f 0.25 (silica gel, pet ether (7)/acetone (3)).




1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[6-aminoheptyl]-3-oxo (6-4)

6-3 (0.44 g) was dissolved in EtOAc (25 ml) cooled to -78° and treated with HCl gas for 5 minutes. The reaction mixture was then stirred at 0° for 30 minutes and the solvent was removed. The residue was purified, by flash chromatography on silica gel eluting with EtOH(9)/H₂O(1)/NH₄OH(1) to provide 6-4 as a white solid. ¹H NMR (300 MHz, CD₃OD) δ 1.42 (4H, m), 1.68 (4H, m), 2.63 (2H, t), 2.88 (2H, t), 3.60 (4H, m), 4.52 (2H, s), 7.60 (1H, d), 7.97 (1H, d), 8.10 (1H, s).


SCHEME 7






10 Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[4-(N-methyl-N-t-butyloxycarbonylamino)butyl]-3-oxo (7-1)

15 Treatment of 1-4 with 4-(N-methyl-N-t-butyloxycarbonylamino)butylamine (prepared as described for 5-3) as described for 1-9 provided crude 7-1. This was purified by flash chromatography on silica gel eluting with EtOAc(7)/hexane(3) to give pure 7-1. R_f 0.3 (silica gel, EtOAc(7)/hexane(3)). ^1H NMR (300 MHz, CDCl_3) δ 1.45 (9H, s), 1.60 (4H, m), 7.52 (1H, d), 8.23 (1H, d), 8.23 (1H, d), 8.50 (1H, s).




25 1H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[4-(N-methyl-N-t-butyloxycarbonylamino)butyl]-3-oxo (7-2)

30 Treatment of 7-1 (1.16 g, 2.08 mmoles) with $\text{LiOH}\cdot\text{H}_2\text{O}$ (0.65 g, 15.5 mmoles) in THF(1)/ CH_3OH (1)/ H_2O (1) (75 ml) as described for 1-10 gave 7-2 as a white solid. ^1H NMR (300 MHz, CD_3OD) δ 1.67 (10H, m), 1.80 (2H, m), 1.89 (2H, m), 3.05 (3H, s), 3.50 (2H, t), 3.88 (2H, t), 4.78 (2H, s), 7.90 (1H, d), 8.45 (1H, d), 8.60 (1H, s).






40 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl)ethyl]-2-[4-(N-t-butyloxycarbonyl-N-methylamino)butyl]-3-oxo (7-3)

45 Treatment of 7-2 (1.04 g, 2.86 mmoles) with β -alanine t-butyl ester (0.54 g, 2.97 mmoles) as described for 1-11 gave crude 7-3. This was purified by flash chromatography on silica gel eluting with hexane(6)/acetone(4) to give 7-3 as an oil. R_f 0.4 (silica gel, EtOAc(7)/hexane(3)). ^1H NMR (300 MHz, CHCl_3) δ 1.46 (18H, m), 1.60 (4H, m), 2.58 (2H, t), 2.83 (3H, s), 3.28 (2H, t), 3.70 (4H, m), 4.45 (2H, s), 7.52 (1H, d), 8.09 (1H, d), 8.11 (1H, s).


7-4

10 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-carboxyethyl]-2-(4-(N-methylamino)butyl]-3-oxo (7-4)

Treatment of 7-3 with HCl gas in EtOAc solution as described for 6-4 gave 7-4 as a white solid.
¹H NMR (300 MHz, CD₃OD) δ 1.67 (4H, m), 2.58 (5H, m), 2.95 (2H, t), 3.50 (4H, m), 4.50 (2H, s), 7.56 (1H, d), 7.97 (1H, d), 8.08 (1H, s).

SCHEME 81-4Et₃N8-11) Boc₂O

2) LiOH

8-2

SCHEME 8 cont'd

5

10

15

20


25

30

35

40

45




Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[(3-aminomethylphenyl) methyl]-3-oxo (8-1)

50

Treatment of 1-4 (2.15 g, 7.49 mmoles) with m-xylenediamine (9.85 mmoles) as described for 1-9 gave crude 8-1. This was purified by flash chromatography on silica gel eluting with CH_3OH (10/ CHCl_3 (NH_4OH) (90) to give pure 8-1 as a white solid. R_f 0.7 silica gel, CH_3OH (10)/ CHCl_3 (NH_4OH) (90).

55




1-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[(3-N-t-butyloxycarbonylaminoethyl)phenyl]methyl]-3-oxo (8-2)

8-1 (1.76 g, 5.67 mmoles) was dissolved in CH_2Cl_2 (25 ml) and treated with Boc_2O (1.50 g, 6.87 mmoles) and Et_3N (6.45 mmoles) as described for 6-2 to give the desired N-protected ester. R_f 0.25 (silica gel, EtOAc (1)/hexane (1)).

^1H NMR (300 MHz, CDCl_3) δ 1.45 (9H, s), 1.65 (1H, m), 2.06 (2H, s), 4.30 (4H, m), 4.81 (2H, s), 7.27 (6H, m), 7.47 (1H, d), 8.22 (1H, d), 8.55 (1H, s).

This acid was treated with $\text{LiOH}\cdot\text{H}_2\text{O}$ as described for 6-2 to provide 8-2 as a white solid. R_f 0.1 (silica gel, CHCl_3 (97)/ CH_3OH (1)/HOAc (1)).


^1H NMR (300 MHz, CD_3OD) δ 1.32 (9H, s), 4.12 (2H, s), 4.38 (2H, s), 4.73 (2H, s), 7.12 (4H, m), 7.25 (1H, m), 7.52 (1H, d).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl)ethyl]-2-[(3-N-t-butyloxycarbonylaminoethyl)phenyl]methyl]-3-oxo (8-3)

Treatment of 8-2 (0.80 g, 2.02 mmoles) with β -alanine t-butyl ester (0.35 g, 2.28 mmoles), BOP (1.35 g, 3.04 mmoles) and Et_3N (14.3 mmoles) as described for 1-11 gave crude 8-3. This was purified by flash chromatography on silica gel eluting with hexane (6)/acetone (4) to give pure 8-3.




^1H NMR (300 MHz, CDCl_3) δ 1.45 (9H, s), 1.47 (9H, s), 2.59 (2H, t), 3.72 (2H, m), 4.30 (4H, s), 4.82 (2H, s), 4.88 (1H, m), 7.28 (5H, m), 7.48 (1H, d), 8.08 (1H, d), 8.19 (1H, s).







1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[(3-aminomethylphenyl)methyl]-3-oxo (8-4)


8-3 (0.872 g, 1.67 mmoles) was dissolved in EtOAc (25 ml) and treated with HCl as described for 6-4 to give pure 8-4.

^1H NMR (300 MHz, CD_3OD) δ 2.58 (2H, t), 3.56 (2H, t), 4.00 (4H, s), 4.42 (2H, s), 7.32 (4H, m), 7.52 (1H, d), 7.95 (1H, d), 8.11 (1H, s).

SCHEME 9.


SCHEME 9 cont'd9-39-49-1Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[(4-amino-1,1,4,4-tetramethyl)butyl]-3-oxo (9-1)

Treatment of 1-4 (2.51 g, 8.74 mmoles) with 1,1,4,4,-tetramethyl-1,4-diaminobutane (1.50 g, 10.40 mmoles) as described for 1-9 provided 9-1. R_f 0.25 silica gel, 10% CH₃OH in CHCl₃/NH₄OH.

9-21-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[(4-N-t-butyloxycarbonylamino)-1,1,4,4-tetramethyl]butyl]-3-oxo (9-2)


9-1 was treated with Boc₂O and Et₃N as described for 6-2 to give the desired Boc-protected ester. R_f 0.3 (silica gel, hexane (7)/acetone/3).

This ester (1.03 g, 2.46 mmoles) was treated with LiOH·H₂O (0.54 g, 12.9 mmoles) in THF (1)/CH₃OH (1)/H₂O (1) (60 ml) as described for 6-2 to give pure 9-2. R_f 0.35 (silica gel, EtOAc).
¹H NMR (300 MHz, CD₃OD) δ 1.10 (6H, s), 1.28 (9H, s), 1.48 (6H, s), 4.60 (2H, s), 7.55 (1H, d), 8.16 (1H, d), 8.26 (1H, s).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-t-butyloxycarbonyl ethyl]-2-[4-(N-t-butyloxycarbonylamino)-(1,1,4,4-tetramethyl)butyl]-3-oxo (9-3)


9-2 (1.05 g, 2.83 mmoles) was treated with β-alanine t-butyl ester (0.48 g, 3.12 mmoles), Et₃N (20.0 mmoles) and BOP (1.91 g, 4.31 mmoles) in CH₃CN (15 ml) as described for 1-11 to provide crude 9-3. This was purified by flash chromatography on silica gel eluting with pet ether (7)/acetone (3) to give pure 9-3. R_f 0.3 silica gel, pet ether (7)/acetone (3).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[(4-amino-1,1,4,4-tetramethyl)butyl]-3-oxo (9-4)

9-3 (1.23 g) was dissolved in EtOAc (25 ml), cooled to -78° and treated with HCl gas as described for 6-4 to give pure 9-4.

¹H NMR (300 MHz, CD₃OD) δ 1.26 (6H, s), 1.53 (8H, m), 2.59 (2H, t), 3.57 (2H, m), 4.63 (2H, s), 7.57 (1H, d), 7.98 (1H, d), 8.06 (1H, s).

SCHEME 10

SCHEME 10 cont'd

5

10

15

20

25


30

35

40

45


50



Methyl-1H-Indole-5-carboxylate, 2,3-dihydro-N-[3-(4-N-t-butyloxycarbonylpiperidinyl)propyl]-3-oxo (10-2)


Treatment of 1-4 (4.59 g, 16.0 mmoles) with 3-(4-N-t-butyloxycarbonylpiperidinyl)propylamine (prepared from 1-6 by nitrile formation followed by catalytic hydrogenation) (4.36 g, 15.6 mmoles) as described for 1-9 gave crude 10-2. This was purified by flash chromatography on silica gel eluting with hexane (3)/ethyl acetate (1) to give pure 10-2.

¹H NMR (300 MHz, CDCl₃) δ 1.10 (2H, m), 1.30 (2H, m), 1.45 (9H, s), 1.68 (4H, m), 2.66 (2H, m), 3.62 (2H, t), 3.95 (3H, s), 4.10 (2H, m), 4.44 (2H, s), 7.52 (1H, d), 8.23 (1H, d), 8.50 (1H, s).

10-31-H-Isoindole-5-carboxylic acid, 2,3-dihydro-N-[3-(4-N-t-butyloxycarbonylpiperidinyl)propyl]-3-oxo (10-3)


Treatment of 10-2 (2.79 g, 6.91 mmoles) with LiOH·H₂O (1.48 g, 35.2 mmoles) in THF (1)/MeOH (1)/ H₂O (1) as described for 1-10 provided 10-3 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 0.95 (2H, m), 1.23 (3H, m), 1.35 (9H, s), 1.66 (3H, m), 2.65 (2H, m), 3.56 (2H, t), 3.96 (2H, bd), 4.50 (2H, s), 7.60 (1H, d), 8.17 (1H, d), 8.30 (1H, s).

10-41-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-(t-butyloxycarbonyl) ethyl]-2-[3-(4-N-t-butyloxycarbonylpiperidinyl)propyl]-3-oxo (10-4)


Treatment of 10-3 (1.28 g, 3.28 mmoles) with β-alanine t-butyl ester (0.64 g, 3.52 mmoles), Et₃N (3.3 mmoles), BOP (2.16 g) in CH₃CN as described for 1-11 gave crude 10-4. This was purified by flash chromatography on silica gel eluting with hexane (7)/ acetone (3) to give pure 10-4.

¹H NMR (300 MHz, CDCl₃) δ 1.09 (2H, m), 1.30 (3H, m), 1.45 (9H, s), 1.68 (4H, m), 2.62 (4H, m), 3.62 (2H, t), 3.70 (2H, t), 4.08 (2H, bd), 4.23 (2H, s), 7.52 (1H, d), 8.10 (1H, d), 8.13 (1H, s).


10-51-H-Isoindole-5-carboxamide, 2,3-dihydro-N-(2-carboxyethyl)-2-[3-(4-piperidinyl)propyl]-3-oxo (10-5)

Treatment of 10-4 (1.18 g) in EtOAc (30 ml) -78° with HCl gas as described for 6-4 gave pure 10-5 as a white solid. R_f 0.4 (silica gel, EtOAc).

¹H NMR (300 MHz, CD₃OD) δ 1.30 (4H, m), 1.67 (4H, m), 1.89 (2H, bd), 2.60 (2H, t), 2.40 (2H, t), 3.19 (2H, bd), 3.58 (4H, m), 4.50 (2H, s), 7.60 (1H, d), 7.99 (1H, d), 8.08 (1H, s).

SCHEME 11


SCHEME 11 cont'd



20

25

HCl (gas)
EtOAc




40

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-methyl-N-2-(carboethoxy)ethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (11-2)

Treatment of 1-10 (0.2 g, 0.54 mmoles) with ethyl 3-(N-methyl)aminopropionate (0.14 g, 1.08 mmoles) (Appl. Polymer Sci., 1969, 13, 227), N-methylmorpholine (1.08 mmoles), and BOP (0.35 g, 0.8 mmoles) in CH₃CN (3 ml) as described for 1-11 gave crude 11-2. This was purified by flash chromatography on silica gel eluting with EtOAc to give pure 11-2 as a white solid.


45

¹H NMR (300 MHz, CD₃OD) δ 1.20 (6H, m), 1.45 (9H, s), 1.67 (2H, q), 1.80 (2H, bd), 2.73 (2H, m), 3.00 (3H, s), 3.08 (1H, bs), 3.71 (2H, t), 3.84 (1H, m), 4.05 (4H, m), 4.17 (1H, m), 4.56 (2H, s), 7.66 (2H, m), 7.77 (1H, s).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-methyl-N-(2-carboxyethyl)]-2-[(2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl)-3-oxo (11-3)]

11-2 (0.23 g, 0.49 mmol) was treated with LiOH·H₂O (0.096 g, 2.3 mmol) as described for 8-2 to give 11-3 as a white solid.






1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-methyl-N-(2-carboxyethyl)]-2-[(4-piperidinyl)ethyl]-3-oxo (11-4)

11-3 (0.2 g, 0.45 mmol) in EtOAc was treated with HCl gas as described for 8-4 to give pure 11-4 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.14 (1H, t), 1.37 (2H, m), 1.50 (1H, m), 1.63 (2H, q), 1.92 (2H, bd), 2.51 (1H, t), 2.67 (1H, t), 2.83 (2H, m), 3.31 (2H, bd), 3.54 (1H, t), 3.60 (2H, t), 3.73 (1H, t), 4.49 (2H, s), 7.57 (2H, q), 7.65 (1H, s).

SCHEME 12



SCHEME 12 cont'dMethyl 3-amino-2,2-dimethylpropionate (12-2)

35 12-1 (Aldrich, 5.0 g, 38 mmoles) in toluene (150 ml) at room temperature was treated with chlorodiphenyl phosphine (49.4 mmoles) followed by imidazole (5.7 g, 83.6 mmoles) and I_2 (12.5 g, 49.4 mmoles) and the resulting brown solution was stirred for 0.5 hours. This mixture was poured into 150 ml saturated Na_2CO_3 solution and the organic layer was separated and washed with saturated Na_2CO_3 solvent, 5% Na_2SO_4 solution, H_2O , and 10% $KHSO_4$ solution. The nearly colorless organic layer was then washed with brine, dried (Na_2SO_4)

40 and the solvent was removed to produce a yellow residue. This was purified by flash chromatography on silica gel eluting with hexane (6)/EtOAc (4) to give the desired iodo intermediate as an oil. R_f 0.9 (silica gel, hexane (6)/EtOAc (4)).


1H NMR (300 MHz, $CDCl_3$) δ 1.38 (6H, s), 3.40 (2H, s), 3.75 (3H, s).

This iodo compound (3.9 g, 16 mmoles) was dissolved in DMSO (80 ml) and treated with NaN_3 (2.1 g, 32 mmoles) at 70° for 2 hours. The cooled reaction next was diluted with EtOAc and extracted with H_2O and brine. The organic phase was washed with brine, dried (Na_2SO_4) and the solvent was removed to give the desired azide as a foam.

1H NMR (300 MHz, $CDCl_3$) δ 1.25 (6H, s), 3.45 (2H, s), 3.75 (3H, s).


This azide (2.0 g, 12.7 mmoles) was dissolved in THF (50 ml) and treated with H_2O (25 ml) and triphenyl phosphine (13.3 g, 50.8 mmoles) at room temperature for 2 hours. The THF was removed under vacuum and the resulting residue was acidified to pH 2-3 with 10% $KHSO_4$ solution. This was filtered to remove triphenyl phosphine and the filtrate was extracted with EtOAc. The acidic aqueous phase was then basified with 10% NaOH and extracted with Et_2O . The combined ether extracts were washed with brine, dried (Na_2SO_4) and the solvent removed to give 12-2 as a clear oil. R_f 0.35 (silica gel, CH_2Cl_2 (9)/ CH_3OH (1)/ H_2O (1)).

55 1H NMR (300 MHz, CD_3OD) δ 1.22 (6H, s), 2.75 (2H, s), 3.75 (3H, s).



10 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[(2-carbomethoxy-2-methyl)propyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (12-3)

Treatment of 1-10 (1.0 g, 2.7 mmoles) with 12-2 (0.524 g, 4.0 mmoles), N-methylmorpholine (4.0 mmoles) and BOP (1.78 g, 4.0 mmoles) in CH₃CN (15 ml) as described for 6-3 provided crude 12-3. This was purified
 15 by flash chromatography, on silica gel eluting with EtOAc (9)/Hexane (1) to give pure 12-3 as a white solid.
¹H NMR (300 MHz, CDCl₃) δ 1.20 (2H, m), 1.33 (6H, s), 1.48 (9H, s), 1.80 (2H, bd), 2.71 (2H, bt), 3.64 (2H, d), 3.73 (2H, t), 3.77 (3H, s), 4.13 (2H, m), 4.44 (2H, s), 6.94 (1H, t), 7.57 (1H, d), 8.11 (1H, d), 8.13 (1H, s).



30 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[(2-carboxy-2-methyl)propyl]-2-[2-(4-piperidinyl)ethyl]-3-oxo (12-4)

12-3 (0.5 g, 1.0 mmoles) was treated with LiOH·H₂O (0.216 g, 5.0 mmoles) as described for 6-2 to give the desired acid as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.13 (2H, m), 1.25 (6H, s), 1.45 (9H, s), 1.65 (2H, m), 1.80 (2H, bd), 2.72 (2H, m), 3.68 (2H, m), 3.70 (2H, t), 4.05 (2H, bd), 4.56 (2H, s), 7.67 (1H, d), 8.04 (1H, dd), 8.15 (s).

This acid (0.40 g) was dissolved in EtOAc and was treated with HCl gas as described for 6-4 to give pure 12-4 as a white solid.

¹H NMR (300 MHz, D₂O) δ 1.14 (6H, s), 1.35 (2H, m), 1.49 (1H, m), 1.60 (2H, q), 1.90 (2H, bd), 2.81 (2H, t), 3.30 (2H, bd), 3.47 (2H, s), 3.57 (2H, t), 4.48 (2H, s), 7.55 (1H, d), 7.82 (1H, d), 7.90 (1H, s).


40


45

50

55

SCHEME 13




SCHEME 13 cont'd

40 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-phenethyl-N-2-carboethoxyethyl]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)ethyl]-3-oxo (13-2)

1-10 (0.388 g, 1.0 mmoles) was treated with ethyl 3-(N-phenethyl)aminopropionate (0.22 g, 1.0 mmoles) (prepared by treatment of phenethylamine with ethyl acrylate), triethylamine (0.243 g, 2.4 mmoles) and BOP (0.53 g, 1.2 mmoles) in DMF (15 ml) and the resulting solution was stirred at room temperature for 18 hours. The solvent was then removed and the residue was diluted with H₂O (100 ml) and extracted with EtOAc (3 x 100 ml portions). The organic phase was washed with 10% KHSO₄ solution, brine, saturated NaHCO₃ solution, brine and dried (Na₂SO₄). Solvent removal gave 13-2 as an oil.


¹H NMR (300 MHz, CDCl₃) δ 1.07-1.35 (6H, m), 1.48 (9H, s), 1.62 (3H, m), 1.75 (2H, bd), 2.72 (4H, m), 3.00 (1H, m), 3.50 (2H, m), 3.67 (2H, t), 3.83 (2H, m), 4.10 (5H, m), 4.38 (2H, s), 6.94 (1H, bs), 7.30 (6H, m), 7.50 (1H, m), 7.67 (1H, m).

13-3

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-phenethyl-N-(2-carboxyethyl)]-2-[2-(4-N-t-butyloxycarbonyl-piperidinyl)ethyl]-3-oxo (13-3)

13-2 (0.60 g, 1.0 mmoles) was treated with LiOH·H₂O (0.127 g, 3.0 mmoles) as described for 6-2 to give 13-3 as a white solid. R_f 0.45 (silica gel, CHCl₃ (9)/MeOH (5)/HOAc (1)).

¹H NMR (300 MHz, CDCl₃) δ 1.17 (2H, m), 1.47 (9H, s), 1.63 (3H, m), 1.75 (2H, bd), 2.67 (2H, t), 2.80 (3H, m), 3.42 (1H, m), 3.57 (1H, m), 3.67 (2H, t), 3.80 (2H, m), 4.08 (3H, m), 4.37 (2H, s), 6.93 (1H, m), 7.25 (6H, m), 7.48 (1H, m), 7.70 (1H, m).

13-4

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[N-phenethyl-N-(2-carboxyethyl)]-2-[2-(4-piperidinyl)ethyl]-3-oxo (13-4)

13-3 was treated with HCl (gas) in EtOAc as described for 6-4 to give pure 13-4 as a white solid. R_f 0.25 (silica gel, EtOH (10)/H₂O (1)/NH₄OH (1)).

¹H NMR (300 MHz, CD₃OD) δ 1.45 (2H, m), 1.62 (2H, m), 1.71 (2H, m), 2.07 (2H, bd), 2.45 (1H, m), 2.78 (2H, m), 2.95 (3H, m), 3.37 (3H, bd), 3.57 (1H, bt), 3.72 (2H, t), 3.83 (2H, m), 3.55 (2H, s), 6.95 (1H, m), 7.20 (4H, bs), 7.33 (1H, bs), 7.45 (1H, bs), 7.55 (1H, m), 7.66 (1H, m).

SCHEME 14.

5

10

15


20

25

30

35

40



45 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[t-butyloxycarbonylmethyl]-2-[3-(4-N-t-butyloxycarbonylpiperidinyl)propyl]-3-oxo (14-1)

Treatment of 10-3 with glycine t-butyl ester as described for 6-3 gave 14-1.


50 ¹H NMR (300 MHz, CDCl₃) δ 1.13 (2H, m), 1.30 (2H, m), 1.41 (9H, s), 1.52 (9H, s), 1.73 (4H, m), 2.69 (2H, t), 3.65 (2H, t), 4.10 (2H, bd), 4.16 (2H, d), 4.45 (2H, s), 7.53 (1H, d), 8.10 (1H, d), 8.22 (1H, s).

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[carboxymethyl]-2-[3-(4-piperidinyl)propyl]-3-oxo (14-2)

Treatment of 14-1 with HCl gas in EtOAc as described for 6-4 gave 14-2 as a white solid.

55 ¹H NMR (300 MHz, CD₃OD) δ 1.30 (4H, m), 1.65 (4H, m), 1.90 (2H, bd), 2.59 (2H, t), 2.90 (2H, t), 3.30 (2H, bd), 3.58 (4H, m), 4.50 (2H, s), 7.58 (1H, d), 7.98 (1H, d), 8.07 (1H, s).

SCHEME 15



Methyl-1H-Isoindole-5-carboxylate, 2,3-dihydro-N-[2-(4-aminobutyl)]-3-oxo(15-1)

1-4(2.56g, 8.92mmoles) was treated with 1,4-diaminobutane (10.9 mmoles) as described for 1-9 to give crude 15-1. This was purified by flash chromatography on silica gel eluting with 25% $\text{CH}_3\text{OH}/\text{CHCl}_3(\text{NH}_3)$ to give pure 15-1 as a solid.


^1H NMR (300 MHz, CD_3OD) δ 1.61 (2H, m), 1.75 (2H, m), 2.90 (2H, t), 3.24 (1H, m), 3.63 (2H, t), 3.85 (3H, s), 4.53 (2H, s), 7.62 (1H, d), 8.18 (1H, d) 8.28 (1H, s).

1-H-Isoindole-5-carboxylic acid-2,3-dihydro-N-[2-(4-N-t-butyloxycarbonylamino)butyl]-3-oxo(15-2)

15-1 (1.11g, 4.24mmoles) was treated with Boc_2O (1.17g, 5.36 mmoles) as described for 3-1. Crude residue was purified by flash chromatography on silica gel eluting with 30% acetone/hexane to give the desired protected ester as an oil. R_f 0.7 silica gel, 30% acetone/hexane.

This ester (0.85g, 2.34mmoles) was dissolved in $\text{THF}(1)/\text{CH}_3\text{OH}(1)/\text{H}_2\text{O}(1)$ (30ml) and treated with $\text{LiOH}\cdot\text{H}_2\text{O}$ (0.52g, 12.4mmoles) as described for 3-2 to give 15-2 as a white solid.


^1H NMR (300 MHz, CD_3OD) δ 1.36 (9H, s), 1.44 (2H, m), 1.66 (4H, m), 3.01 (2H, t), 3.60 (2H, t), 4.54 (2H, s), 7.62 (1H, d), 8.20 (1H, d), 8.35 (1H, s).

15-3

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-(t-butyloxycarbonyl)ethyl]-2-[4-(N-t-butyloxycarbonyl)butyl]-3-oxo(15-3)

Treatment of 15-2 (0.75g, 2.07mmoles) in CH₃CN (12ml) with β-alanine t-butyl ester (0.39g, 2.54mmoles), Et₃N (14.3 mmoles) and BOP (1.40g, 3.16 mmoles) as described for 3-3 gave crude 15-3. This was purified by flash chromatography on silica gel eluting with 75% EtOAc/hexane to give pure 15-3 as a white solid. R_f 0.25 (silica gel, 75% EtOAc/hexanes).


¹H NMR (300 MHz, CDCl₃) δ 1.42 (9H, s), 1.44 (9H, s), 1.52 (2H, m), 1.77 (2H, m), 2.55 (2H, t), 3.19 (2H, m), 3.67 (4H, m), 4.43 (2H, s), 7.00 (1H, bt), 7.52 (1H, d), 8.09 (1H, d), 8.10 (1H, s).

15-4

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[2-carboxyethyl]-2-[4-aminobutyl]-3-oxo(15-4)

Treatment of 15-3 (0.51g, 1.07mmoles) in EtOAc with HCl gas as described for 3-4 provided pure 15-4 as a white solid.

¹H NMR (300 MHz, D₂O), δ 1.63 (4H, m), 2.64 (2H, t), 2.92 (2H, t), 3.52 (4H, m), 4.46 (2H, s), 7.55 (1H, d), 7.81 (1H, d), 7.85 (1H, s).




1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[ethyl-3-(2(S)-aminopropionate)]-2-[2-(4-N-t-butyloxycarbonyl-piperidiny)]-3-oxo (16-2)

A solution of **1-10** (1.5g, 3.87 mmoles) in DMF (15ml) at room temperature was treated with carbonyl diimidazole (0.627g, 3.87 mmoles) (CDI) and after 2 hours this solution was added dropwise to a DMF solution of ethyl 2(S),3-diaminopropionate (1.5g, 7.74 mmoles) and N-methylmorpholine (23.2 mmoles). The reaction mixture was then stirred at room temperature for 16 hrs.


The solvent was then removed and the residue was dissolved in EtOAc and 10% aqueous KHSO_4 solution. The aqueous phase was separated, washed with EtOAc and made basic to pH 12. This was extracted with EtOAc, and the extracts were combined, washed with brine, and dried (Na_2SO_4). Solvent removal provided **16-2**.

^1H NMR (300 MHz, CD_3OD) δ 1.24 (2H, m), 1.46 (3H, t), 1.43 (9H, s), 1.66 (2H, q), 1.80 (2H, bd), 3.67 (4H, m), 4.10 (2H, bd), 4.17 (2H, q), 4.57 (2H, s), 7.04 (1H, d), 7.67 (1H, m), 8.06 (1H, m), 8.17 (1H, d).



10 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-[2(S)-aminopropanoic acid]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)-3-oxo]-3-oxo (16-3)

Treatment of 16-2 (0.6 g, 1.2 mmoles) with LiOH·H₂O (0.25 g, 6.0 mmoles) as described for 1-10 gave 16-3.
 15 ¹H NMR (300 MHz, D₂O) δ 0.92 (2H, m), 1.27 (9H, s), 1.46 (4H, m), 2.58 (2H, t), 3.48 (4H, m), 3.83 (2H, bd), 4.38 (2H, s), 6.96 (1H, s), 7.50 (1H, d), 7.82 (1H, d), 7.87 (1H, s).




25

1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-[2(S)-methylsulfonylamino]propanoic acid]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)-3-oxo]-3-oxo (16-6)

30 A solution of 16-6 (0.55 g, 1.2 mmoles) in H₂O (15ml)/dioxane (3ml) was cooled to 0-10° and treated with 1N NaOH soln. (1.5ml) and methane sulfonyl chloride (2.4 mmoles) in 3 ml dioxane was added dropwise while also adding 1N NaOH solution to keep the pH at 10-12. This cycle of CH₃SO₂Cl addition at basic pH was carried out 5 times at which point all 16-6 was consumed. The acidity was carefully adjusted to pH 2-3 with 10% KHSO₄ solution and this was extracted with EtOAc (4 portions). The combined organics were washed with brine, dried
 35 (Na₂SO₄) and the solvent removed. The residue was purified by flash chromatography on silica gel eluting with CH₂Cl₂ (9)/MeOH (0.8)/HOAc (0.8) to give 16-6 as a white solid. R_f 0.31.
¹H NMR (300 MHz, CD₃OD) δ 1.25 (2H, m), 1.45 (9H, s), 1.65 (2H, q), 1.80 (2H, bd), 2.72 (2H, m), 2.97 (3H, s), 3.70 (3H, m), 3.86 (1H, m), 4.05 (2H, bd), 4.34 (1H, m), 4.56 (2H, s), 7.66 (1H, d), 8.08 (1H, d), 8.19 (1H, s).


40



45

50 1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-(2(S)-methylsulfonylamino)propionic acid]-2-[2-(4-piperidyl)ethyl]-3-oxo (16-7)

Treatment of 16-6 (0.22 g, 0.39 mmoles) with HCl gas in EtOAc as described for 1-12 gave 16-7 as a white solid.
 55 ¹H NMR (300 MHz, D₂O) δ 1.35 (2H, m), 1.59 (2H, m), 1.87 (2H, bd), 2.78 (2H, bt), 2.95 (3H, m), 3.27 (2H, bd), 3.55 (3H, m), 3.78 (1H, m), 4.20 (1H, m), 4.48 (2H, s), 7.56 (1H, m), 7.87 (1H, m), 7.95 (1H, bs).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-(2(S)-n-butylsulfonylamino)propanoic acid]-2-[2-(4-N-t-butyloxycarbonylpiperidinyl)]-3-oxo (16-8)

Treatment of 16-3 (0.836 mmoles) with n-butylsulfonyl chloride (1.67 mmoles) as described for 16-6 gave 16-8 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 0.85 (6H, m), 1.13 (2H, m), 1.35 (4H, m), 1.45 (9H, s), 1.65 (2H, m), 1.75 (2H, m), 2.70 (2H, m), 3.04 (2H, t), 3.68 (2H, m), 3.83 (1H, m), 4.04 (2H, bd), 4.53 (2H, s), 7.62 (1H, d), 8.05 (1H, d), 8.18 (1H, s).



1-H-Isoindole-5-carboxamide, 2,3-dihydro-N-[3-(2(S)-n-butylsulfonylamino)propionic acid]-2-[2-(4-piperidinylethyl)]-3-oxo (16-9)

Treatment of 7-8 in EtOAc with HCl gas as described for 1-12 gave pure 16-9 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 0.59 (2H, t), 1.12 (2H, m), 1.35 (2H, m), 1.50 (2H, m), 1.59 (2H, m), 1.90 (2H, bd), 2.80 (2H, t), 2.98 (2H, t), 3.29 (2H, bd), 3.42 (1H, m), 3.60 (2H, t), 3.70 (1H, m), 4.50 (2H, s), 7.59 (1H, d), 7.91 (1H, d), 7.98 (1H, s).

Claims

1. A fibrinogen receptor antagonist of the following formula:



wherein:

A, B, C and D independently represent a carbon atom or a nitrogen atom;



E is $-(\text{CHR}^1)_m-(\text{CHR}^2)_n-\text{F}-(\text{CHR}^3)_o-(\text{CHR}^4)_p-$;

$-(\text{CHR}^1)_m-\text{CR}^2=\text{CR}^3-(\text{CHR}^4)_n-\text{F}-$; or

$-\text{F}-(\text{CHR}^1)_m-\text{CR}^2=\text{CR}^3-(\text{CHR}^4)_n-$,



wherein

m, n, o, and p are integers chosen from 0-2; and F is an optional substituent, which when present is chosen from:



or $-\text{NR}^1\text{R}^2$;

X is $-\text{NR}^1\text{R}^2$,



or a 4- to 10- membered mono- or polycyclic aromatic or nonaromatic ring system containing 0, 1, 2, 3 or 4 heteroatoms selected from N, O and S and either unsubstituted or substituted with R^1 , R^2 , R^3 or R^4 , wherein R^1 , R^2 , R^3 and R^4 are independently selected from the group consisting of hydrogen,


C_{1-10} alkyl,
 aryl C_{0-8} alkyl,
 oxo,
 thio,
 5 amino C_{0-8} alkyl, C_{1-3} acylamino C_{0-8} alkyl,
 C_{1-6} alkylamino C_{0-8} alkyl,
 C_{1-6} dialkylamino C_{0-8} alkyl,
 C_{1-4} alkoxy C_{0-8} alkyl,
 10 carboxy C_{0-8} alkyl, C_{1-3} alkoxycarbonyl C_{0-8} alkyl,
 carboxy C_{0-8} alkyloxy and
 hydroxy C_{0-8} alkyl;

Y is C_{0-8} alkyl,
 15 C_{0-8} alkyl- NR^3 -CO- C_{0-8} alkyl,
 C_{0-8} alkyl-CONR³- C_{0-8} alkyl,
 C_{0-8} alkyl-O- C_{0-8} alkyl,
 C_{0-8} alkyl-S(O_n)- C_{0-8} alkyl, or
 C_{0-8} alkyl-SO₂-NR³- C_{0-8} alkyl-,
 20 C_{0-8} alkyl-NR³-SO₂- C_{0-8} alkyl-,
 C_{1-8} alkyl-CO- C_{0-8} alkyl;


Z is



30 O, S, SO, SO₂, SO₂(CH₂)_m, (CH₂)_mSO₂, (CH₂)_m,



SO₂NR³,



45 NR³SO₂ or CR³=CR⁴,
 wherein m is 0-6;

R⁵ is hydrogen
 C_{1-8} alkyl,
 50 C_{0-8} alkylcarboxy C_{0-8} alkyl,
 C_{0-8} alkyloxy C_{0-8} alkyl,
 hydroxy C_{0-8} alkyl,
 aryl C_{0-8} alkyl, or
 halogen;


55 R⁶ is hydrogen,
 C_{1-8} alkyl,
 aryl C_{0-8} alkyl,

C₃₋₈ cycloalkyl C₀₋₆ alkyl,
 C₀₋₆ alkylcarboxy C₀₋₆ alkyl, carboxy C₀₋₆ alkyl,
 C₁₋₄ alkyloxy C₀₋₆ alkyl,
 hydroxy C₀₋₆ alkyl, provided that any of which groups may be substituted or
 unsubstituted independently with R¹ or R², and provided that, when two R⁶ groups are attached
 to the same carbon, they may be the same or different;

R⁷ is hydrogen, fluorine
 C₁₋₈ alkyl,
 C₃₋₈ cycloalkyl,
 aryl C₀₋₆ alkyl,
 C₀₋₆ alkylamino C₀₋₆ alkyl,
 C₀₋₆ dialkylamino C₀₋₆ alkyl,
 C₁₋₈ alkylsulfonylamino C₀₋₆ alkyl,
 aryl C₀₋₆ alkylsulfonylamino C₀₋₆ alkyl,
 C₁₋₈ alkyloxycarbonylamino C₀₋₆ alkyl,
 aryl C₀₋₆ alkyloxycarbonylamino C₀₋₆ alkyl,
 C₁₋₈ alkylcarbonylamino C₀₋₆ alkyl,
 aryl C₀₋₆ alkylcarbonylamino C₀₋₆ alkyl,
 C₀₋₈ alkylaminocarbonylamino C₀₋₆ alkyl,
 aryl C₀₋₈ alkylaminocarbonylamino C₀₋₆ alkyl,
 C₁₋₈ alkylsulfonyl C₀₋₆ alkyl,
 aryl C₀₋₈ alkylsulfonyl C₀₋₆ alkyl,
 C₁₋₈ alkylcarbonyl C₀₋₆ alkyl,
 aryl C₀₋₈ alkylcarbonyl C₀₋₆ alkyl,
 C₁₋₈ alkylthiocarbonylamino C₀₋₆ alkyl,
 aryl C₀₋₈ alkylthiocarbonylamino C₀₋₆ alkyl
 wherein groups may be unsubstituted or substituted with one or more substituents selected
 from R¹ and R², and provided that when two R⁷ groups are attached to the same carbon atom,
 they may be the same or different;

R⁸ is hydroxy,
 C₁₋₈ alkyloxy,
 aryl C₀₋₆ alkyloxy,
 C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyloxy,
 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyloxy, or
 an L- or D-amino acid joined by an amide linkage and wherein the carboxylic acid moiety of
 said amino acid is as the free acid or is esterified by C₁₋₆ alkyl.

2. A compound of Claim 1, having the formula





II wherein:

E is: $-(CHR^1)_m-(CHR^2)_n-F-(CHR^3)_o-(CHR^4)_p-$,
 $-(CHR^1)_m-CR^2=CR^3-(CHR^4)_n-F-$, or
 $-F-(CHR^1)_m-CR^2=CR^3-(CHR^4)_n-$,
 where m, n, o and p are integers 0-2.


F is an optional substituent which when present is chosen from:

O; -CR¹R²;



or -NR¹R²; and X, Y, R¹, R², R³, R⁴, R⁶, R⁷ and R⁸ are as previously defined in Claim 1.

3. A compound of Claim 2, having the formula:



wherein:

E is: -(CHR¹)_m-F-(CHR²)_n-,

-CR¹=CR²-F-, or



-F-CR¹=CR²-,

where m and n are integers 0-2

and

F is an optional substituent which when present is chosen from:

O; -CR¹R²;



or -NR¹R²-

X is

-NR¹R² or a 4- to 10-membered mono- or polycyclic aromatic or non-aromatic ring system containing 0, 1 or 2 heteroatoms chosen from N or O and either unsubstituted or substituted with R¹ and R², wherein

R¹ and R² are independently chosen from:

hydrogen,

C₁₋₆ alkyl,

aryl C₀₋₆ alkyl,

carboxy C₀₋₆ alkyl,

hydroxy C₀₋₆ alkyl,

C₁₋₃ alkyloxy C₀₋₆ alkyl, or

amino C₀₋₆ alkyl;

Y is

C₀₋₆ alkyl,






C₁₋₆ alkyl-CO-C₀₋₆ alkyl, or

C₀₋₆ alkyl-NR³-CO-C₀₋₆ alkyl;


R⁹ and R⁷ are as previously defined in Claim 1, and
R⁸ is

hydroxy,
C₁₋₆ alkyloxy,
aryl C₁₋₄ alkyloxy, or
C₁₋₄ alkylcarbonyloxy C₁₋₄ alkyloxy.


4. A compound of Claim 1 selected from the group of



5




10




15

20




25

30




35

40








45


50



55




5




10

15




20

25




30



35

40




45

and

50

55



5. A compound as claimed in any of Claims 1 to 4 for use in therapy.

6. A pharmaceutical composition comprising a compound as claimed in any of Claims 1 to 4 and a pharmaceutically acceptable carrier.
7. A composition as claimed in Claim 6 which also comprises a thrombolytic agent.
- 5 8. The composition of Claim 7 wherein the thrombolytic agent is a plasminogen activator or streptokinase.
9. A composition as claimed in Claim 6 which also comprises an anticoagulant.
- 10 10. The composition of Claim 9, wherein the anticoagulant is heparin or warfarin.
11. A composition as claimed in Claim 6 which also comprises an antiplatelet agent.
12. The composition of Claim 11, wherein the antiplatelet agent is aspirin.
- 15 13. The use of a compound as claimed in any of Claims 1 to 4 for the manufacture of a medicament for inhibiting the binding of fibrinogen to blood platelets.
14. The use of a compound as claimed in any of Claims 1 to 4 for the manufacture of a medicament for inhibiting the aggregation of blood platelets.
- 20 15. The use of a compound as claimed in any of Claims 1 to 4 for the manufacture of a medicament for the prevention or treatment of thrombus or embolus formation.
- 25 16. A composition comprising a compound as claimed in any of Claims 1 to 4 in combination with two or more agents selected from a thrombolytic agent, an anticoagulant agent, and an antiplatelet agent and a pharmaceutically acceptable carrier.
17. The composition of Claim 16, wherein the thrombolytic agent is a plasminogen activator or streptokinase, the anticoagulant agent is heparin or warfarin, and the antiplatelet agent is aspirin.

30

35

40

45

50

55



**European Patent
Office**

EUROPEAN SEARCH REPORT

Application Number

EP 92 30 9924

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|---|---|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. Cl.5) |
| A | US-A-4 010 274 (GIRALDI) 1 March 1977 | 1-17 | C07D09/46 A61K31/395 |
| A | --- | 1 | |
| A | US-A-3 322 631 (SPRAGUE) 30 May 1967 compounds in column 1, lines 15-25 | 1-17 | |
| A | EP-A-0 332 528 (RHONE-POULENC SANTE) 13 September 1989 compounds I | 1-17 | |
| A | FR-A-2 612 185 (FARMITALIA CARLO ERBA) 16 September 1988 compounds I | 1-17 | |
| The present search report has been drawn up for all claims | | | TECHNICAL FIELDS SEARCHED (Int. Cl.5) |
| | | | C07D A61K |
| Place of search MUNICH | | Date of completion of the search 21 JANUARY 1993 | Examiner GETTINS M. P. |
| CATEGORY OF CITED DOCUMENTS | | T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document | |
| X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document | | | |